

BAND SPECTRUM OF THALLIUM IODIDE

By P. TIRUVENGANNA RAO AND K. R. RAO

(Received for publication, Nov. 4, 1948)

Plates VA and VB

ABSTRACT. An extensive band system attributed to the diatomic molecule thallium iodide, has been obtained for the first time in emission between $\lambda 5300$ – $\lambda 3800$ in a high frequency discharge with external electrodes. The bands at the violet end are red-degraded; those in the central portion are headless; at the red end they are diffuse and broad. A number of them occur in well-separated groups, each consisting of four or five equi-spaced components with an interval of about 28 units; the intensity among these rises gradually to a maximum and then falls off. The system has been interpreted as due to the transition ${}^3\Pi_1 - {}^1\Sigma^+$ with the origin near the extreme violet end. A vibrational analysis which includes most of the band heads, has led to the following formula:—

$$\nu = 25794.2 + (91.2 u' - 0.09 u'^2) - (121.0 u'' - 0.09 u''^2)$$

A second brief system of about 25 slightly violet degraded bands is also obtained between $\lambda 3680$ – $\lambda 3600$. Four diffuse groups of bands, overlaid by a continuum, are measured. The system has been interpreted as the component ${}^3\Pi_0^+ - {}^1\Sigma^+$ having the same lower state as that of the main system. Electron transitions are compared for the observed band systems of the four halide molecules of thallium.

INTRODUCTION

As part of a detailed investigation into the molecular spectra of a series of heavy diatomic fluorides, Howell (1937) studied the thallium fluoride spectrum and reported several points of unusual interest. Extending this investigation to the other halides of the same element, thallium chloride and thallium bromide, Howell and Coulson (1938, 1941) were enabled to make a comparison of the electronic levels and transitions for these three molecules. The band spectrum of the remaining halide, thallium iodide, does not appear to have been analysed.

In 1929, Butkow recorded a few bands in absorption by a column of thallium iodide vapour, but no vibrational scheme was reported; nor was any work published on the emission spectrum of the molecule, as far as the authors are aware. In the course of a systematic investigation in our laboratory of the band spectra of diatomic molecules and particularly, the iodides, the spectrum of thallium iodide has been photographed and it is the purpose of the present paper to describe this spectrum and to report the analysis of it as interpreted by the authors.

EXPERIMENTAL

The discharge tube employed is a simple straight one of pyrex glass about 30 cm. long and 1 cm. in diameter, drawn out at one end into an adapter,

the other end being closed by a quartz window. The substance, a pure sample of thallium iodide obtained from B.D.H. is spread inside the tube over a length of about 8 to 10 cms. The tube is continuously evacuated by a Cenco Hyvac pump through a row of absorbing and condensing towers. The discharge is excited by a low power H.F. oscillator constructed for the purpose and described elsewhere (Ramasastry, 1947) in detail, working at a frequency of 5 to 10 megacycles and using external copper electrodes about 5 cm. apart. The substance was heated at regular intervals for a steady maintenance of the discharge; such intermittent heating was found necessary also for suppressing the iodine bands which would otherwise obscure the thallium iodide system. The discharge is intense green, due to the thallium radiation $\lambda 5350$.

The bands were obtained also in a discharge from a $1/4$ Kw. transformer in an ordinary H type tube containing the vapour. The H.F. discharge was found more favourable for the production of the system.

A number of spectrographs were used to photograph the spectrum—a Fuess glass instrument, Hilger large glass and quartz Littrow instruments and a 10-foot concave grating in the first order. On account of the peculiar nature of the spectrum, measurements had to be made on the small as well as the large dispersion spectrograms on a number of plates for different regions. The usual international iron arc standards were adopted to reduce the plates.

DESCRIPTION OF THE SPECTRUM

The spectrum cannot be adequately described without reference to the photographs, some of which are reproduced in Plates VA and VB. Figures *a*, *b* and *c* of Plate VA are glass Fuess spectrograms from $\lambda 4900$ to $\lambda 3800$, giving a general view of the entire system of bands. The development of the system below $\lambda 3900$, which is poor in the glass instrument, is shown in the quartz Littrow picture, Fig. *d*, Plate VA. The heads give the appearance of a complex system, with one marked regularity. Over a wide range towards the violet end, they occur in well-separated groups each consisting of about four or five equi-spaced components with an interval of approximately 28 wave-number units. The fluctuations of intensity show a kind of wave appearance such as is observed in the band systems of sulphur or of selenium. There are altogether about 200 band heads within the region $\lambda 5300$ to $\lambda 3800$, being very crowded in the central portion. The bands near the red-end show greater diffuseness and many of them appear double. In each of the groups at the violet end, the intensity rises and falls suggesting the formation of short sequences. At the violet end, the heads are sharper and clearly red-degraded; diffuseness sets in and increases as one goes towards the red. The peculiarities of appearance noticed by Howell in the thallium chloride and thallium fluoride bands are not observed in the iodide system—namely the

doubling at the head and the occurrence of the so called "absorption lines" in the emission spectrum, nor does the iodide system resemble closely that of the bromide, which consists of long progressions with very intense, sharp and line-like heads near the shortwave side.

The kind of evidence of the existence of isotope effect that is found in thallium chloride and thallium bromide bands is absent in this system as iodine has no isotopes, leading to a special difficulty in the interpretation of its structure. The effect due to the thallium isotopes, however, is seen in the doubling or broadening of the heads in the region of longer wave lengths. An evidence of this is seen in Plate VB Figs. *e* and *f*, which are enlargements of the spectra in this region taken by the Fuess and the concave grating respectively. Fig. *g*, in Plate VB shows another system between $\lambda_{3680-3600}$ which is also ascribed to the thallium iodide molecule. It consists of three or perhaps four continuous patches superposed by distinct sequences of bands having nearly the same wave number interval of about 28 units as in the first system.

RESULTS


Table I is a catalogue of the band heads assigned to the first system and Table II gives the data for the second system. The tables are self-explanatory.

TABLE I

Wave length	Int.	Wave number	Classifica- tion v', v''	Wave length	Int.	Wave number	Classifica- tion v', v''
3812.96	2	26218.9	5,0	23.93	4	25477.5	2,4
17.52	2	26187.6	6,1	28.33	4	25448.9	3,5
21.64	2	26159.4	7,2	32.95	3	25419.0	4,6
				37.76	2	25388.0	5,7
25.60	2	26132.3	4,0				
29.96	2	26102.6	5,1	3940.90	3	25367.8	.
34.60	2	26071.0	6,2	45.30	1	25339.5	
				49.80	1	25310.6	
38.60	2	26043.8	3,0	51.92	1	25297.0	
42.90	2	26014.7	4,1				
47.40	2	25984.2	5,2	56.51	3	25267.7	5,8
				60.86	4	25239.9	6,9
51.90	2	25953.9	2,0	65.73	3	25208.9	7,10
56.66	2	25921.9	3,1	69.74	4	25183.5	8,11
61.41	2	25890.0	4,2	81.91	1	25106.5	
				84.26	2	25091.7	
65.31	2	25863.8	1,0				
69.43	2	25836.3	2,1	88.99	3	25061.9	8,12
73.75	3	25807.5	3,2	93.37	4	25034.5	9,13
				96.80	4	25012.4	10,14
77.88	5	25780.0	0,0	4005.96	1	24955.8	
88.28	3	25751.8	1,1	10.04	1	24930.4	
				13.82	1	24906.9	
96.80	2	25654.8	0,1				
3900.78	5	25628.7	1,2	17.00	4	24887.2	10,15
04.92	4	25601.5	2,3	21.81	4	24857.4	11,16
09.19	3	25573.5	3,4	26.53	3	24828.3	12,17
17.84	1	25517.1		31.49	2	24797.7	13,18
20.10	1	25502.4		44.52	1	24730.1	

TABLE I (contd.)

Wave length	Int.	Wave number	Classification v', v''	Wave length	Int.	Wave number	Classification v', v''
46.30	1	24707.0	12,18	4306.18	3	23215.9	28,43
50.33	3	24682.4	13,19	11.23	3	23188.7	29,44
55.09	3	24653.4	14,20	15.92	5	23163.5	30,45
57.88	4	24636.5	15,21	21.32	4	23134.6	31,46
60.43	1	24621.0					
62.57	1	246.8.0					
64.91	1	24593.9	16,22	4326.14	3	23108.8	32,47
				31.62	3	23079.6	33,48
79.48	2	24506.0	15,22	37.43	2	23048.7	34,49
84.31	4	24477.1	16,23				
88.80	5	24450.2	17,24	42.58	5	23021.3	31,47
93.88	2	24419.8	18,25	47.69	3	22994.8	32,48
				52.88	3	22966.9	33,49
4113.18	5	24305.3	18,26	57.87	3	22940.6	34,50
18.23	4	24275.5	19,27	62.61	2	22915.6	35,51
23.38	3	24245.1	20,28	67.24	1	22891.4	36,52
27.99	1	24218.1					
28.13	1	24217.2	21,29	69.33	4	22880.4	32,49
				74.91	3	22851.2	33,50
39.76	2	24149.2	19,28	80.09	2	22924.2	34,51
42.75	3	24131.8	20,29				
45.21	1	24117.5		85.04	3	22758.4	31,49
47.59	3	24103.6	21,30	89.84	4	22773.5	32,50
				95.61	2	22743.6	33,51
4152.24	3	24076.6	22,31	4400.93	2	22716.1	34,52
57.14	2	24048.2	23,32				
62.37	1	24018.0	24,33	06.75	2	22686.1	31,50
				11.98	3	22659.2	32,51
67.02	0	23991.2	21,31	17.47	4	22631.1	33,52
71.35	1	23966.3	22,32	22.57	5	22605.0	34,53
77.64	1	23930.2	23,33	28.89	2	22572.7	35,54
82.50	3	23902.1	24,34	33.53	2	22549.1	36,55
87.27	4	23875.2	25,35	39.63	1	22518.1	37, 6
92.36	2	23846.2	26,36				
96.36	1	23823.5		44.56	1	22493.1	34,54
				50.28	4	22464.2	35,55
4207.73	2	23759.1	25,36	55.70	3	22436.9	36,56
10.09	1	23745.8		60.82	2	22411.0	37,57
13.28	5	23727.8	26,38	66.40	1	22383.8	38,58
17.36	4	23704.9	27,38	67.52	1	22377.5	
				72.64	1	22351.9	39,59
23.26	1	23671.8	24,36	75.01	1	22340.1	
27.60	1	23647.9	25,37				
32.69	2	23619.0	26,38	77.88	5	22325.8	36,57
36.15	1	23590.7		84.19	3	22294.3	37,58
38.27	4	23587.9	27,39	88.87	3	22271.0	38,59
40.78	2	23574.0		94.57	3	22242.9	39,60
43.61	3	23558.2	28,40				
48.45	3	23531.4	29,41	4500.35	1	22214.3	36,58
53.41	3	23504.0	30,42	06.62	3	22183.4	37,59
				11.64	5	22158.7	38,60
58.31	3	23476.9	27,40	17.43	4	22130.3	39,61
63.47	3	23448.5	28,41	22.88	3	22103.6	40,62
69.66	5	23414.5	29,42	28.54	2	22076.0	41,63
74.76	2	23386.6	30,43	34.16	2	22048.6	42,64
77.41	1	23372.1					
79.75	1	23359.3	31,44	39.76	2	22021.4	39,62
81.95	1	23347.3		45.57	3	21993.3	40,63
84.33	3	23334.4	32,45	47.86	1	21982.2	
				51.71	3	21963.6	41,64
89.26	4	23307.5	29,43	57.14	5	21937.5	42,65
94.34	4	23279.9	30,44	62.79	4	21910.3	43,66
99.80	3	23250.4	31,45				






Fig.

Fig. f.




Fig. g.

Thallium iodide spectrum (Fe arc comparison)

Band Spectrum of Thallium Iodide

189

TABLE I (contd.)

Wave length	Int.	Wave number	Classification <i>v', v''</i>	Wave length	Int.	Wave number	Classification <i>v', v''</i>
73.54	3	21858.0	41.65	4728	2	21143	
76.98	1	21842.4		4737	4	21103	51,79
80.18	3	21827.1	42.66	4749	2	21052	48,77
86.07	4	21799.1	43.67	4785	2	20894	
91.27	5	21774.4	44.68	4795	3	20849	49,80
93.30	3	21764.8		4822	3	20733	49,81
4615.15	2	21661.7	44.69	4832	1	20692	
20.85	5	21635.2	45.70	4843	1	20642	48,81
27.99	2	21601.6	46.71	4853	1	20601	
				4898	2	20411	48,83
32.30	2	21581.5	43.69	4932	1	20269	
38.73	2	21551.6	44.70	4947	3	20211	47,84
43.85	2	21527.9	45.71	4968	2	20122	46,84
50.66	2	21496.3	46.72	4981	3	20070	
56.21	3	21470.7	47.73	5038	2	19844	48,88
62.19	2	21443.2	48.74	5060	1	19760	47,88
67.78	2	21417.5	49.75	5059	1	19723	48,89
				5077	2	19690	49,90
75.40	4	21382.6	46.73	5091	1	19636	47,89
85.89	3	21334.7	48.75	5113	0	19554	
91.94	2	21307.2	49.76	5118	1	19532	
97.05	1	21284.0	50.77	5150	2	19412	47,91
				5134	2	19396	
4703.05	4	21256.9	47.75	5186	2	19277	48,93
10.02	4	21225.4	48.76	5235	0	19098	
22.10	4	21171.1	50.78	5255	1	19025	49,96

TABLE II

Wave length	Int.	Wave number	Classification
3588.87	0		
84.98	1	27100.9	(0,0)
81.24	1	27129.5	(1,1)
77.86	1	27157.1	(2,2)
74.12	1	27182.0	(3,3)
70.79	2	27209.7	(4,4)
67.07	1	27234.4	(5,5)
63.80	1	27262.0	(6,6)
60.59	1	27286.3	(7,7)
57.40	1	27310.3	(8,8)
53.94	1	27334.1	(9,9)
50.31	1	27360.0	(10,10)
		27387.2	(11,11)
44.61	1		
41.04	2	27430.0	(8,7)
37.45	1	27456.9	(9,8)
34.51	1	27484.0	(10,9)
30.12	1	27506.2	(11,10)
26.38	1	27539.5	(12,11)
		27567.9	(13,12)
19.65	1		
16.30	1	27619.1	(11,9)
12.68	1	27644.7	(12,10)
08.91	1	27672.4	(13,11)
05.61	0	27701.3	(14,12)
		27726.7	(15,13)
3588.03	1		
85.73	1	27862.5	(16,13)
		27885.1	(17,14)

VIBRATIONAL ANALYSIS

The well marked regularity in the band system mentioned earlier served as a starting point in the analysis. The clearly separated groups, each consisting of a few almost equi-spaced bands with gradual rise and fall off in intensity, are considered as different sequences. Two chief difficulties, however, presented themselves in working out the analysis further; (1) The location of the origin of the system. At this stage a comparison is made with the known systems of the other thallium halides whose structures are well-established. The main system in each of these molecules occurs in the near ultraviolet with the system origin located at the most refrangible end; if a similar feature is expected in the main system of thallium iodide and if intensity considerations and variations of the interval between the successive groups are also taken into account, it seemed that the strongest band at $\nu=25780.0 \text{ cm.}^{-1}$ may be the (0,0) band of the system. (2) The second difficulty is in respect of the magnitude of the values of ω' and ω'' ; several alternatives of choice are available; the interval of 28 cm.^{-1} between the nearly equi-spaced bands in each sequence suggested the approximate value of the difference between ω' and ω'' . The individual values may be one of the following sets:—

90	120
120	150
150	180
180	210

Petrikaln and Hochberg (1933) have quoted (p. 229, table 14.) a value 150 for ω'' . It is a common observation in band spectra that the vibrational constants are not much altered in the case of a row of heavy diatomic molecules, formed by a successive addition of an electron. The following table illustrates this feature in the halides of Au, Hg, Tl etc.

TABLE III
Ground State Constants

	Au	Hg	Tl	Pb	Bi
F	—	490.8	475.0	506.9	512.0
Cl	383	293.4	287.5	304.2	308.4
Br	—	186.3	192.1	207.5	209.5
I	—	125.7	121.0	—	163.9

TABLE IV

Vibrational Constants of Thallium Halides *

	ν_e	ω_e'	$x_e'\omega_e'$	ω_e''	$x_e''\omega_e''$
TlI [†]	35180.7	439.8	8.60	175.0	1.89
TlCl	31054.2	216.9	6.80	287.5	1.24
TlBr	29191.5	108.3	5.15	192.1	0.39
TlI	25685.9	91.0	0.20	121.0	0.18

Table III suggests that for thallium iodide, the value of ω'' may be approximately either 150 cm.^{-1} or 120 cm.^{-1} . A further comparison of the ground and excited state constants of the thallium halides as shown in Table IV indicates that of these two probable values the latter *i.e.* 120 cm.^{-1} is probably to be preferred, as it would simultaneously lead to an ω' value which is quite consistent with the corresponding values of the other halides. If 150 cm.^{-1} be adopted for ω'' , the value of $\omega' \dagger$ will be 120 which is entirely out of step. Hence, of the different possible alternative schemes of the vibrational analysis of the bands, the authors have adopted the one which gives $\omega' \sim 90 \text{ cm.}$ and $\omega'' \sim 120$ units.

The classification of the bands arrived on the above basis is shown in the last column of Table I. It has been possible to include all the bands right up to $\lambda 5300$ into a single vibrational scheme, although, as will be seen later, a doubt exists regarding the classification of the bands in the extreme red.

VIBRATIONAL CONSTANTS AND ISOTOPE EFFECT

Confirmation of the vibrational analysis from a study of the isotope effect is desirable in dealing with systems of this type. But the difficulty has been that iodine has no isotopes. Thallium has two isotopes of mass 205 and 203 in the abundance ratio 2.4:1 and it should be possible to detect the weaker 203 isotopic head for the $\text{Tl}^{203}\text{I}^{127}$ molecule; particularly when band heads with very large vibrational quantum numbers are identified. For this purpose the vibrational constants for the system are derived as accurately as possible. The source of error, here arises from the absence of long progressions or sequences and the very low value of the anharmonic constant, apart from the uncertainty in the measurement of wavelengths due to diffuseness. A graphical method is made use of, trying to average out the discrepancies in the $\Delta G(v)$ values for both upper and lower states. Curves

* Values for TlI, contained in this table, are those obtained in the present work.

† A value of 30 may also be possible, but is equally out of step.

in Fig. 1 show the manner in which $\Delta G(v')$ and $\Delta G(v'')$ vary. If linear curves are drawn through the points, the estimated constants are

$$\begin{aligned}\omega_e' &= 91.2 & \omega_e'' &= 121.2 \\ x_e' \omega_e' &= 0.10 & x_e'' \omega_e'' &= 0.09 \\ \nu_e &= 25794.7.\end{aligned}$$




FIG. 1




FIG. 2

The ν_e values calculated from bands spread-over the entire region of the spectrum seldom differ by more than 5 wave number units. Hence the usual quantum formula with the above vibrational constants represents the entire system.

Using the above approximate values of the vibrational constants and the factor $(\rho - 1) = 0.0019$ for the thallium iodide molecule, the isotopic separations given by the usual formula (Jevons' Report, p.213)

$$\nu^i - \nu = (\rho - 1) (\omega_e' u' - \omega_e'' u'') - (\rho^2 - 1) (x_e' \omega_e' u'^2 - x_e'' \omega_e'' u''^2) +$$

is calculated for a number of bands. The order of separation for bands at the extreme red is about six wave number units; *e.g.*, for the (39,58) and (49,73) bands it is 5.7 and 6.8 wave number units respectively. Plate VB, Fig. *f* is a concave grating spectrogram of the bands in the red region taken in order to identify the $Tl^{203}I^{127}$ isotopic heads. It is easily seen that the bands are extremely diffuse and unsuitable for measurement. But evidence is found in these bands either of a broadening or in several cases of the splitting of the head into two components with an estimated interval of the same order as the expected isotopic displacement. No further detailed study of the isotope effect has been possible in this system.




FIG. 3

Energy levels of TlF, TlCl, TlBr and TlI.

INTENSITY DISTRIBUTION

The intensity distribution among the bands, if the proposed analysis is correct, is an exceptional feature of the system. The distribution is represented in a vibrational matrix in Table V in which the visual estimates of intensity are shown in place of the wave numbers. A prominent difference between this and the other halide systems of thallium is noticeable in respect of this intensity distribution. Predissociation is detected in thallium fluoride, chloride and bromide bands; the upper state being cut off at a specific vibrational level which is different for the different molecules. This predissociation is considered as probably responsible for an identical intensity distribution among the bands both in emission and absorption. In thallium bromide in particular, there are just six progressions which are observed up to

a very high value of ν'' . No evidence of predissociation exists in the thallium iodide system. Table V shows that bands occur involving high values of both ν' and ν'' . This peculiar distribution may in part, arise from the absence of predissociation and may also be due to the low values of xw for the ground and excited states and the relative magnitude of ω' and ω'' . These latter constants are neither nearly equal which would lead to a narrow Condon parabola nor is their ratio too large or too small which would give a very wide and open parabola. The ratio is such that just a few bands alone may appear in a sequence or a progression.

If a curve is drawn through the more intense bands, we obtain, instead of the usual parabola, one in which one arm is exceptionally long. At the end of the system, the intensity distribution is erratic. It must be stated that due to this and the possible large errors in measurement, the classification of the bands in this region may be uncertain.

One view which prevailed from the outset owing to the exceptional intensity distribution was (Rao and Rao, 1948) to regard the system as made up of two different overlapping systems. The crowding of the bands and the rise in general intensity in the central portion of the system at about $\lambda 4300$ (Refer Plate VA Fig. a) are suggestive of such a division. Just at this region, the origin of the second system may be situated with the system extending towards the red. But a more detailed examination revealed that at the red end of the system, the order of the intervals between the groups is about 110 cm.^{-1} . If these bands form part of a second system with the origin near $\lambda 4300$, the value of 110 is too low, since the anharmonic constant is expected to be small, while, if they form part of an extensive system with the origin at $\lambda 3878$, the value of 110 is of the right order of magnitude. It is because of this consideration and also of the difficulty, as discussed later, in interpreting the electronic transition of a brief system of bands observed on the violet end, that the view of the division of the main system into two components had to be abandoned. All the bands from $\lambda 3800$ to $\lambda 5300$ are regarded by the authors as a single system. (Rao and Rao, 1948).

ELECTRONIC STATES

Miescher and Wehrli (1934 and 1935), Howell and others (1937) discussed the electronic states of the halides of group III elements Al, Ga, etc., in relation to the observed band spectra. Holst (1934) and Jennergren (1948) have established the ground states of the molecules AlCl and AlBr as ${}^1\Sigma^+$. For the other molecules as well the ground state may be in all probability a similar ${}^1\Sigma^+$. Two systems are known in most of these molecules one of which is a singlet lying in the far-ultraviolet and the other a doublet system occurring in the near ultraviolet. Both have a common lower state ${}^1\Sigma^+$ arising from the electron configuration $\sigma^2\pi^4$. The excitation of one the π electrons produces the configuration $\sigma^2\pi^3\sigma$ giving rise to the molecular states ${}^1\pi$, ${}^3\pi_0^+$, ${}^3\pi_0'$, ${}^3\pi_1$ and ${}^3\pi_2$ (inverted). As shown by Howell for thallium flouride chloride and

TABLE VI
Vibrational analysis of $^3\Pi_g^+ - ^1\Sigma^+$ system of thallium iodide

v'' v'	0	1	2	3	4	5	6	8	9	10	11	12	13	14	$\Delta G''(v')$
0	27100.9														
1		27129.5													143.7
2			27157.1												146.6
3				27182.0											149.9
4					27209.7										140.7
5						27234.4									145.4
6							27262.0								132.9
7								27286.3							133.4
8									27310.3						132.4
9										27334.					135.8
10											27360.0				
11										27387.2					
12											27419.1				
13												27444.7			
14													27472.4		
15														27507.9	
16															27701.3
17															27833.7
															27725.7
															27862.5
															27885.1
$\Delta G''(v'')$								119.7	122.8	118.5	112.1	104.5	107.0		

bromide, the singlet system may be identified as $\sigma^2\pi^3\sigma, {}^1\pi-\sigma^2\pi^4, {}^4\Sigma^+$ and the doublet system as a forbidden transition ${}^3\pi_0^+-{}^1\Sigma^+$ and ${}^3\pi_1-{}^1\Sigma^+$. The singlet system is predominant in the aluminium, gallium and indium halides and the doublet system is either absent or poorly developed, while for TlI, the two systems are of equal intensity. Such a relative intensity behaviour is consistent with the above assignment of transitions and indicates, as Howell remarked, that the binding from Ga→In→Tl approaches more and more to Hund's case (c) as the molecule gets heavier.* In the heaviest molecules of this type, namely TlI, one has therefore to anticipate the predominance of the forbidden transitions ${}^3\pi_0-{}^1\Sigma^+$ and ${}^3\pi_1-{}^1\Sigma^+$. On this account, it was at first considered probable that the extensive system observed for TlI in the appropriate wave length region might consist of two overlapping components.

Further scrutiny of the plates has revealed a second short region between $\lambda 3680-\lambda 3600$ in which distinct band heads, overlaid by a continuum, are observed. These consist of four groups of nearly equi-spaced components with an interval of about 28 units. The four groups appear to be different sequences, the strongest of them at $\lambda 3670$ perhaps being the $\Delta v=0$ sequence. The interval of 120 cm.^{-1} also exists between certain heads in these groups which is of the same order as that of the lower state of the main system. There is an impression of a degradation of the bands in the direction of shorter wave lengths. With the aid of these regularities, the vibrational scheme in Table VI has been set up which includes all the band heads. Wave length and other data of these bands are given in Table II. The following approximate constants have been determined from the analysis, and curve in Fig 2.

$$v_{(0,0)} = 27101; \quad \omega_e' \sim 151; \quad \omega_e'' \sim 120.$$

If the main system is treated as itself comprising of the two components ${}^3\pi_0-{}^1\Sigma^+$ and ${}^3\pi_1-{}^1\Sigma^+$, the interpretation of the brief system becomes difficult. On the other hand when the main system, in spite of its unusual intensity characteristics, is regarded as a single component due to the transition ${}^3\pi_1-{}^1\Sigma^+$, the brief system may be the other component ${}^3\pi_0-{}^1\Sigma^+$. Such an interpretation would give the ${}^3\pi$ interval as 1321 cm.^{-1} , which is quite in keeping with the corresponding intervals in the other halides as shown in Table VII.

Further, the characteristic features of this brief system are intrinsically such as to support this view. While for example in TlI both the components are observed consisting of band heads, in TlCl and TlBr this second component may presumably be a repulsive predissociating state. The diffuseness of the bands overlaid by a continuum and the fragmentary development of the

* When the coupling is of this type, the states ${}^3\pi_0^+$ and ${}^3\pi_1$ should be denoted as ${}^3_0^+$ and 3_1 .

system may hence be consistent with the assignment of the system as ${}^3\pi_0^+ - {}^1\Sigma^+$.

TABLE VII

	Al	Ga	In	Tl
F	—	—	—	1682
Cl	—	332	796	(1107)*
Br	—	370	786	(872)*
I	199	330	649	1321

* Approximate estimates from the continua.

A schematic diagram of the electronic transitions for the halides of thallium is given in Fig. 3 which is an extension of those presented by Howell.

Bands due to the transition ${}^1\pi - {}^1\Sigma$, if they are present in TII, are yet to be discovered. They are not found in emission under the conditions in which they are photographed in this work. Absorption experiments too, which are in progress, may not be expected to reveal this system.

ANDHRA UNIVERSITY,
WALTAIR.

REFERENCES

- Butkow, 1929, *Zell. f. Physik.*, **58**, 232.
 Holst, 1934, " " **93**, 65.
 Howell, 1937, *Proc. Roy. Soc.*, **160**, 242.
 Howell and Coulson, 1938, *Proc. Roy. Soc.*, **166**, 238.
 " " 1941, *Proc. of Phy. Soc.*, **53**, 706.
 Jennergren Carl Gustav, 1948, *Nature*, **161**, 315
 Miescher and Wehrli, 1934 *Helv. Phy. Acta.*, **73**, 298.
 " " , 1935, " " **8**, 279.
 Petrikalu & Hochberg, 1933, *Zeit. f. Physik*, **86**, 214.
 Ramasastry, 1947, *Ind. Jour. Phys.*, **21**, 272.
 Rao Tiruvengauna & Rao, K. R., 1948, *Curr. Sci.*, **17**, 121.
 " " **17**, 182.