

REFERENCES

- Bloor J.E., Schlubitz J., Walden C. C. & (in part) Demerdache A. 1964 *Canad. J. Chem.* **42**, 2201
 Gurnovich G P., Sevechenko A N. & Solv'ev K N. 1963 *Soviet Phys. Usp (English Translation)* **6**, 173.
 Sidorov A. N & Korlyar I P. 1961 *Opt. Spectry. U.S.S.R. (English Translation)* **XI**, 175.

Indian J. Phys. **44**, 660-663, (1970)

Crystal and molecular structure of *p*-dimethylamino-benzaldehyde hydrobromide

BY J. K. DATTA GUPTA and N. N. SAHA

Crystallography and Molecular Biology Division,

Saha Institute of Nuclear Physics, Calcutta-9

(Received 2 March 1971)

p-Dimethylaminobenzaldehyde ($C_9H_{11}NO$) is a biologically important compound. Among its various biological functions mention may be made of its special role in differentiating between true scarlet fever and serum eruptions. The structural analysis of this compound in the form of its different hydrohalides and metal complexes has been undertaken by us with an ultimate view to correlate the structural features with biological functions. This short communication deals with the crystal structure of *p*-dimethylaminobenzaldehyde hydrobromide.

The compound was prepared in our laboratory by treating *p*-dimethylaminobenzaldehyde with 30% HBr. Single crystals were grown by slow evaporation of an aqueous solution of this compound between 35°-40°C. The crystals thus grown are needle shaped, *c*-axis being parallel to needle axis. As the crystal was found to be unstable under normal atmospheric condition, it was placed in a sealed thin walled glass capillary while taking X-ray photographs. The crystals belong to monoclinic space group $P2_1/c$ having unit cell dimensions

$$a = 12.65 \text{ \AA}$$

$$b = 10.20 \text{ \AA}$$

$$c = 7.42 \text{ \AA}$$

$$\beta = 90^\circ 30'$$

Density data ($\rho_m = 1.54 \text{ gm/cc}$, $\rho_c = 1.59 \text{ gm/cc}$) indicate that there are four formula units of $(C_9H_{11}NO)HBr$ per unit cell. Multiple-film equi-inclination Weissenberg technique was used to collect three-dimensional intensity data for layers

$hk0$ to $hk5$ and $h0l$. The positions of four heavy atoms (bromine) in the unit cell were located from Patterson synthesis projected along c - and b -axes. Three dimensional Fourier synthesis was computed with phases of the heavy atom on CDC 3600 using the program written by Blount. A spoke and bead model was constructed at this stage satisfying the stereochemistry

Refinement was carried out by the full matrix least squares method using isotropic temperature factors. Program used was the modified (Srikanta) one of




Figure 1. *p*-Dimethylaminobenzaldehyde hydrobromide molecule viewed along b -axis.




Figure 2. Intermolecular packing projected along c^* -axis. o' and o'' refer to the oxygen atoms of the molecules above and below the paper respectively.

Busing, Martin & Levy (1962). A few cycles of refinement brought the discrepancy factor $R = \frac{\sum ||F_o - |F_c||}{\sum |F_o|}$ down to 0.16. Further refinement was made using anisotropic temperature factors for the non-hydrogen atoms and the R value at this stage was 0.11. A difference Fourier synthesis computed at this stage revealed the positions of hydrogen atoms. Taking the contribution of hydrogen atoms, the R value was 0.10. Atomic parameters and the intramolecular bond lengths are given in tables 1 and 2, respectively. The molecule as viewed down b -axis has been shown in figure 1. Amino nitrogen atom has tetragonal configuration and is bonded to the benzene ring, two methyl groups and the bromine atom. The bond between the amino nitrogen and bromine atom is a hydrogen bond of the type N-H...Br, length being 3.13 Å. Least squares and best plane calculations show that the angle between the plane containing dimethyl group $-\text{N} \begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix}$ and the plane of the benzene ring is 61°, whereas the aldehyde group $-\text{C} \begin{matrix} \text{O} \\ \text{H} \end{matrix}$ is almost coplanar with the benzene ring. In a similar structure of Wurster's red bromide (Tanaka *et al* 1968) the dimethyl-amino group is perfectly coplanar with the benzene ring. The molecules are held together in space by a number of Van der Waals bonds between carbon atoms and oxygen atom of adjacent molecules. Three dimensional packing of the molecules is shown in figure 2. A paper in details will be published elsewhere shortly.

TABLE 1. Atomic Parameters
a) Fractional co-ordinates

Atom	x/a	y/b	z/c
Br	0.16113	0.02253	0.31758
O	0.62371	0.22132	0.99875
N	0.17272	0.02105	0.73858
C(1)	0.59546	0.14497	0.88826
C(2)	0.48982	0.11714	0.85915
C(3)	0.45969	0.03155	0.72883
C(4)	0.35603	-0.00447	0.68238
C(5)	0.27976	0.05755	0.78086
C(6)	0.30310	0.15014	0.90431
C(7)	0.41073	0.17619	0.94685
C(8)	0.09388	0.12191	0.80171
C(9)	0.15305	-0.11201	0.78936

(b) Anisotropic temperature coefficients*

Atom	β_{11}	β_{22}	β_{33}	β_{12}	β_{13}	β_{23}
Br	0.00482	0.01012	0.01088	-0.00057	-0.00034	-0.00048
O	0.00504	0.00928	0.01968	-0.00289	0.00249	-0.00131
N	0.00064	0.00159	0.00769	0.00048	0.00041	-0.00087
C(1)	0.00192	0.01304	0.00665	0.00056	-0.00456	0.00264
C(2)	0.00431	0.00601	0.01805	0.00048	0.00050	0.00286
C(3)	0.00205	0.01087	0.01146	0.00192	0.00095	0.00900
C(4)	0.00504	0.00719	0.00843	-0.00013	0.00065	0.00280
C(5)	0.00377	0.00428	0.01632	0.00076	0.00151	-0.00509
C(6)	0.00035	0.00954	0.00627	-0.00149	-0.00042	-0.00026
C(7)	0.00511	0.00902	0.00610	0.00029	0.00462	0.00310
C(8)	0.00944	0.00888	0.05073	-0.00384	-0.00033	0.01191
C(9)	0.00828	0.01257	0.01663	0.00444	0.00205	0.00896

*In the expression $T^{-1} = \exp[-(\beta_{11}h^2 + \beta_{22}k^2 + \beta_{33}l^2 + 2\beta_{12}hk + 2\beta_{13}hl + 2\beta_{23}kl)]$

TABLE 2. Intramolecular bond lengths and bond angles

Bond length (Å)		Bond angles (degrees)	
Br—N	3.127	Br—N—C(5)	105
O—C(1)	1.184	Br—N—C(8)	106
N—C(5)	1.436	Br—N—C(9)	105
N—C(8)	1.510	C(8)—N—C(9)	116
N—C(9)	1.431	C(8)—N—C(5)	112
C(1)—C(2)	1.381	C(9)—N—C(5)	110
C(2)—C(3)	1.355	C(6)—C(5)—N	122
C(2)—C(7)	1.341	C(4)—C(5)—N	116
C(3)—C(1)	1.402	C(1)—C(5)—C(6)	122
C(4)—C(5)	1.370	C(7)—C(6)—C(5)	119
C(5)—C(6)	1.347	C(3)—C(4)—C(5)	114
C(6)—C(7)	1.420	C(2)—C(7)—C(6)	121
		C(2)—C(3)—C(4)	126
		C(7)—C(2)—C(3)	115
		C(1)—C(2)—C(3)	120
		C(1)—C(2)—C(7)	123
		O—C(1)—C(2)	121

REFERENCES

- Husung W. R., Martin K. O. & Levy H. A. 1962 *Least Squares Refinement (XPLS) Program*
 Tanaka J. & Sakabe, N. 1968 *Acta Cryst.* **B24**, 1315