

throws light on the relative stability of the bonds, such variation shows that strength of the chemical bond is in the order  $M-F > M-Cl > M-Br$ . The difference in electronegativities between central atom and ligand atom also supports this result. Such information has also been reported by Sanyal & Dixit (1973) employing stretching force constant  $f_r$ . The interaction constant,  $F$ , follows the same pattern as the stretching force constant  $K$  while the angle bending constant  $D$  shows opposite trend of variation. The other interaction constants do not follow regular pattern.

#### ACKNOWLEDGMENTS

The authors are grateful to Prof. Dr. A. Muller, University of Dortmund, West Germany for his continued interest and encouragement. They are also thankful to Dr. P. Sharan, Principal, Meerut College, Meerut for providing necessary facilities in the department.

#### REFERENCES

- Krasser W. 1969 Schwingungsspektren und Kraft Konstanten gefarbtter anorganischer verdindungen, 104.  
 Pandey A. N., Sharma D. K. & Dublish A. K. 1973 *Spectroscopy Letters* **6**, 491.  
 Ramaswamy K. & Muthusubramanian P. 1971 *Ind. J. Phys.* **45**, 477.  
 Sanyal N. K. & Dixit L. 1973 *Ind. J. Pure and Appl. Phys.* **11**, 452.  
 Sharma D. K., Pandey A. N. & Goel R. K. 1974 *Ind. J. Pure and Appl. Phys.* **12**, 455.  
 Wilson E. B. Jr., Decius D. C. & Cross P. C. 1955 *Molecular vibrations* (McGraw Hill Book Co. Inc., New York).

*Indian J. Phys.* **48**, 854-856 (1974)

## On the infrared absorption spectrum of 4-chloro-2-fluorotoluene molecule

G. JOSHI AND K. K. SHARMA

*Department of Physics and Mathematics, Indian School of Mines, Dhanbad*

(Received 27 February 1974)

The vibrational spectra of toluene and its monohalogenated derivatives have been reported by Pitzer & Scott (1943), Wilmshurst & Bernstein (1957), Mooney (1964), Joshi & Singh (1966, 1967). Deb & Banerjee (1960) have reported the spectra of some di-chlorotoluenes in different states. Recently, Singh & Thakur (1969) have reported the vibrational spectra of some di-fluorotoluenes. But no work seems to have been reported on the vibrational spectra of substituted toluenes where two hydrogen atoms of the phenyl ring have been replaced by

two different halogen atoms. The present communication reports the vibrational frequencies and the analysis of one such molecule viz., 4-chloro-2-fluoro-toluene in liquid state.

The infrared spectrum of the compound was recorded in liquid state of Perkin-Elmer spectrophotometer Model 521 using different cell lengths. The spectrum lies in the region 300–3500  $\text{cm}^{-1}$ .

If we assume  $\text{CH}_3$  group to behave as a point mass and lie in the plane of the molecule then 4-chloro-2-fluorotoluene molecule may belong to  $C_s$  point group.

Table 1. Infrared absorption spectrum of 4-chloro-2-fluorotoluene

*Fundamental vibrations*

Infrared spectrum $\text{cm}^{-1}$	Intensity	Assignment
3076	16	$a'$ , C-H stretching
3037	15	$a'$ , C-H stretching
2960	20	$a'$ , C-H asym. stretching (in $\text{CH}_3$ group)
2930	27	$a'$ , C-H asym. stretching (in $\text{CH}_3$ group)
2865	16	$a'$ , C-H sym. stretching
1613	60	$a'$ , C=C stretching
1587	80	$a'$ , C=C stretching
1490	100	$a'$ , C=C stretching
1445	35	$a'$ , C-H asym. bending (in $\text{CH}_3$ group)
1381	30	$a'$ , C-H sym. bending (in $\text{CH}_3$ group)
1290	40	$a'$ , C-H in-plane bending
1266	62	$a'$ , C-H in-plane bending
1225	80	$a'$ , C-F stretching
1187	63	$a'$ , C- $\text{CH}_3$ stretching
1123	52	$a'$ , C-H in-plane bending
1075	100	$a'$ , $\text{CH}_3$ rocking
1037	15	$a'$ , C-C-C in-plane bending
992	30	$a'$ , C-C-C in-plane bending
890	90	$a'$ , C=C stretching
855	70	$a''$ , C-H out-of-plane bending
833	14	$a''$ , C-H out-of-plane bending
805	70	$a''$ , C-H out-of-plane bending
792	42	$a'$ , C-F in plane bending
752	22	$a'$ , C-Cl stretching
575	45	$a'$ , C- $\text{CH}_3$ in-plane bending
460	11	$a''$ , C-F out-of-plane bending
443	35	$a'$ , C-C-C in-plane bending
370	23	$a'$ , C-Cl in-plane bending
305	11	$a''$ , C- $\text{CH}_3$ out-of-plane bending

Out of the 39 fundamental vibrations, 30 will belong to the vibrations associated with the ring and rest to  $\text{CH}_3$  group. Considering the Z-axis as the two fold axis in the plane of the molecule, the 30 fundamental vibrations get divided into the respective species as  $(21a' - 9a'')$  for the  $\text{C}_6$  group. The assignments have been made mostly on the basis of their intensities and positions in the spectrum. Further guidance has been taken from the vibrational assignments of molecules containing the methyl group.

Frequencies observed corresponding to the fundamental vibrations along with their intensities and probable assignments have been given in table 1.

#### ACKNOWLEDGMENT

Author is thankful to Prof. S. N. Mitra, Head of the Department of Physics and Mathematics and to Director, Indian School of Mines, Dhanbad for their encouragement during the course of this work. Author's sincere thanks are due to Prof. P. Venkateswarlu, Indian Institute of Technology, Kanpur for providing the facilities for recording the spectra on Perkin-Elmer infrared spectrophotometer 521.

#### REFERENCES

- Deb K. K. & Banerjee S. B. 1960 *Ind. J. Phys.* **34**, 554.  
Joshi G. & Singh N. L. 1966 *Sp. Chim. Acta.* **22**, 1501.  
Joshi G. & Singh N. L. 1967 *Sp. Chim. Acta.* **23**, 1341.  
Mooney E. F. 1964 *Sp. Chim. Acta.* **20**, 1343.  
Pitzer K. S. & Scott D. W. 1943 *J. Am. Chem. Soc.* **65**, 803.  
Singh V. B. & Thakur G. 1969 *Ind. J. Pure and Applied Phys.* **6**, 107.  
Wilmshurst J. K. & Bernstein H. J. 1957 *Can. J. Chem.* **35**, 911.

*Indian J. Phys.* **48**, 856-858, (1974)

## An X-ray crystallographic study of p-aminophenylarsonic (arsanilic) acid

A. CHATTERJEE AND S. P. SEN GUPTA

*Department of General Physics & X-rays,  
Indian Association for the Cultivation of Science, Calcutta 700032*

(Received 19 March 1974)

Since very little structural work has so far been made with the important organic analytical reagent arsonic acid and its derivatives which are commonly used in the detection and determination of heavier elements namely, zirconium, scandium, bismuth, tantalum, thorium, etc. (Welcher 1948), systematic studies