

## **<sup>35</sup>Cl Nuclear quadrupole resonance in some chloromethylanilines**

S K Roy, B Amitha and J Uchil

Materials Science Division, Mangalore University,  
Mangalagangothri-574 199, India

Received 16 November 1990, accepted 25 April 1991

**Abstract :** <sup>35</sup>Cl Nuclear quadrupole resonances (NQR) have been detected in 4-chloro-2-methyl, 3-chloro-2-methyl and 5-chloro-2-methylaniline. 4-Chloro-2-methylaniline is found to occur in two polymorphic phases  $\alpha$  and  $\beta$ . The temperature variation of NQR frequencies of 3-chloro-2-methyl and  $\alpha$  and  $\beta$  phases of 4-chloro-2-methylaniline have been studied in the temperature range from 77 K upto the respective melting points. The reliability of Hammett's  $\sigma$  parameter is also discussed.

**Key words :** NQR, polymorphism, temperature dependence, Hammett's  $\sigma$  value.

**PACS No. :** 76.60. Gv

### **1. Introduction**

Nuclear quadrupole resonance in chloromethylaniline had been observed for the first time by Biedenkapp and Weiss (1968) in 4-chloro-2-methylaniline at 77 K and the reported frequency is 34.134 MHz. Reports on other members of this family are not available in literature. Presently, four isomeric compounds of chloromethylaniline including 4-chloro-2-methylaniline have been subjected to NQR investigation and resonances are observed in 4-chloro-2-methyl (m.p. = 300 K), 3-chloro-2-methyl (m.p. = 275 K) and 5-chloro-2-methylaniline (m.p. = 295 K). However, NQR line could not be detected in 2-chloro-6-methylaniline (m.p. = 275 K). The temperature dependence of NQR frequencies and the polymorphism observed in 4-chloro-2-methylaniline are reported in this paper.

### **2. Experimental**

The analar grade chemicals used in this investigation were obtained from M/S Fluka, Switzerland. All these compounds are liquids at room temperature (301 K) and therefore further purification was done by distillation. The samples were solidified by cooling it slowly so as to get moulded in glass vials and used as sample tubes.

A frequency modulated self-quenched super-regenerative oscillator (SRO)-detector of Dean's type (Das and Hahn 1958) coupled with a lock-in-amplifier and

a strip-chart recorder were used to detect the signals. A block diagram of the experimental setup is shown in Figure 1. The SRO generates a family of side bands at intervals of quench frequency corresponding to NQR absorption. However, the strength of the side bands goes on diminishing symmetrically as one moves away from the centre. One of the signals in 4-chloro-2-methylaniline was strong enough to be displayed on the cathode ray oscilloscope. A moderately high modulation depth of about 6-8 KHz was employed throughout to detect the weak signals. The resonance frequency was fixed using the standard beat method employing a very stable signal generator. The accuracy of frequency measurement was within  $\pm 1$  KHz. The temperature of the sample was measured within an accuracy of  $\pm 0.5$  K by making use of a copper-constantan thermocouple. The NQR





Figure 1. Block diagram of the experimental setup.

spectrum at 77 K in each compound was recorded by housing the sample coil in a hollow conical shaped metallic conductor attached to the end of the probe and by immersing it in a liquid nitrogen bath. The temperature variation work was carried out by allowing the sample to warm up slowly as the liquid nitrogen was allowed to evaporate out naturally.

### 3. Results and discussion

The NQR frequency obtained from the freshly solidified 4-chloro-2-methylaniline at 77 K is found to be 33.766 MHz. This shows a decrease of 368 KHz from the reported value (Biedenkapp and Weiss 1968). However, it is found that there is a shift in NQR frequency if this sample is stored at temperatures well below the melting point (300 K) of the material, for a long period. By monitoring the NQR frequency regularly, at every 24 hours after preparation of the sample, it has been found that the intensity of the NQR signal decreases gradually. After about 21 days, a completely new signal instead of the previous one starts growing. The signal strength of this NQR line become stabilized within a week and there is no change thereafter either in frequency or in intensity. The increase in the frequency is 370 KHz and it is nearly the same as the reported frequency (Biedenkapp and Weiss 1968). These observations reveal that there exist polymorphism in 4-chloro-

2-methylaniline. These observations can be explained through a model similar to the one used to explain polymorphism in monochloroacetic acid (McDonald and Hacobian 1982). At room temperature 4-chloro-2-methylaniline is a liquid and it is taken to ice temperature in order to get it solidified. As the process of solidification is fast enough, it is similar to quenching and the metastable state (termed as  $\alpha$ -phase hereafter) of this solid phase is obtained in presence of lattice strains and a number of possible lattice defects. As the strain is released with time, the movement of the defects takes place leading to the averaging of electric field gradient (EFG) and hence the decrease in intensity of the signal. Finally, a stable state with no strain but a number of static defects may lead to a state of equilibrium. This solid phase is termed as  $\beta$ -phase. The NQR spectra obtained from both  $\alpha$  and  $\beta$ -phases of 4-chloro-2-methylaniline at a quench frequency of about 30-35 KHz are given in Figure 2. The signal strength of the signal in the




**Figure 2.** The characteristics of typical NQR spectra obtained in  $\alpha$  and  $\beta$ -phase of 4-chloro-2-methylaniline with the identical recording sensitivity.

$\beta$ -phase is found to be quite high compared to that of  $\alpha$ -phase. The variation of NQR frequencies of both the phases from 77 K upto the onset of melting are given in Figure 3. The smooth variation of NQR frequencies suggest that there is no further transition either in  $\alpha$  or in  $\beta$ -phases over this temperature range.

NQR signals are also detected and reported here for the first time in 3-chloro-2-methyl and 5-chloro-2-methylaniline. No polymorphism is found to exist in these two compounds. A single signal with frequency 33.714 MHz is obtained in 3-chloro-2-methylaniline at 77 K. The variation of NQR frequency with temperature, shown in Figure 3, suggest that there is no phase transition present in this compound. In the case of 5-chloro-2-methylaniline, the recorded NQR spectra showed two weak signals of signal-to-noise ratio (S/N) around three near the melting point (293 K). The measured frequencies are 33.417 and 32.998 MHz. As the signal strengths are too low, temperature dependence of NQR frequencies could not be carried out. At 77 K the lower frequency signal, although weak, could be detected and fixed without difficulty. However, the higher frequency

signal was smeared out and it was not possible to fix the frequency with certainty. The NQR frequencies and line widths at 77 and 273 K and the S/N of all the lines are given in Table 1. As the recorded signal in 5-chloro-2-methylaniline is very weak a proper line width measurement was not possible.



**Figure 3.** Temperature dependence of NQR lines in the two compounds of chloromethylaniline.  $\circ\text{---}\circ\text{---}\circ\text{---}\circ$  and  $\circ\text{---}\circ\text{---}\circ\text{---}\circ$  represent  $\alpha$  and  $\beta$ -phases of 4-chloro-2-methylaniline, respectively and  $\Delta\text{---}\Delta\text{---}\Delta\text{---}\Delta$  represent 3-chloro-2-methylaniline. Continuous dashed lines represent the estimated frequencies in 4-chloro-2-methylaniline.

Fourth degree polynomials fitted to the observed frequency variation with temperature by the method of least squares using an IBM compatible computer for the two compounds are as follows :

#### 1. 4-Chloro-2-methylaniline

$$\alpha\text{-phase } \nu(T) = 33,878 - 1.2173T - 0.2670 \times 10^{-3}T^2 - 0.4265 \times 10^{-5}T^3 - 0.4028 \times 10^{-8}T^4 \quad (\text{r.m.s.d.} = 5.45 \text{ KHz})$$

$$\beta\text{-Phase } \nu(T) = 34,248 - 1.3807T - 0.2749 \times 10^{-3}T^2 - 0.3112 \times 10^{-5}T^3 + 0.2382 \times 10^{-8}T^4 \quad (\text{r.m.s.d.} = 4.75 \text{ KHz})$$

Table I. <sup>13</sup>Cl NQR frequencies in some chloromethylanilines.

Compound	$\nu$ (MHz)		$\nu_0$ (MHz)		Line width (KHz)		S/N
	77 K	273 K	Computed	Extrapolated	77 K	273 K	
1. 4-Chloro-2-methylaniline $\alpha$ -phase	33.766	33.240	33.878	33.880	9	8.5	15 (77 K)
	34.116	33.627	34.248	34.245	7	7.5	23 (77 K)
	34.116*	33.721*	34.182*				
2. 3-Chloro-2-methylaniline	33.714	33.221	33.813	33.820	7	8	12 (77 K)
	34.975*	34.592*	35.047*				
3. 5-Chloro-2-methylaniline	—	33.490	—	—	—	—	3 (273 K)
	33.688	33.065	—	—	—	—	3 (273 K)
	34.513*	34.101*	34.582*				

\*Estimated frequencies using eq. (1).

## 2. 3-Chloro-2-methylaniline

$$\nu(T) = 33,813 - 1.1645T - 0.2354 \times 10^{-3}T^2 - 0.3764 \times 10^{-5}T^3 - 0.4657 \times 10^{-8}T^4$$

(r.m.s.d. = 4.73 KHz)

where r.m.s.d. is the root mean square deviation of least squares fit and  $\nu(T)$  is the NQR frequency in KHz at a temperature  $T(K)$ . The NQR frequencies at OK ( $\nu_0$ ) obtained by the computer fitting and the ones extrapolated from the experimental curves are also given in Table 1.

The NQR frequencies are also estimated using the following equation proposed by Setty and Uchil (1984)

$$\begin{aligned} \nu(\text{in MHz}) = & \left( 34.915 + 1.005 \sum_i \sigma_i \right) - \left( 0.57 + 0.05 \sum_i \sigma_i \right) 10^{-3}T \\ & - \left( 0.46 + 0.05 \sum_i \sigma_i \right) 10^{-5}T^2 \end{aligned} \quad (1)$$

where the summation extends over all substituents at ortho, meta and para positions relative to the  $^{35}\text{Cl}$  nucleus of interest.

The Hammett's  $\sigma$ -values made use of in the calculation are  $o\text{-CH}_3 = 0.292$  (Bray and Barnes 1957),  $m\text{-CH}_3 = -0.069$ ,  $p\text{-CH}_3 = -0.170$ ,  $m\text{-NH}_2 = -0.161$  and  $p\text{-NH}_2 = -0.660$  (Biedenkapp and Weiss 1968). The estimated frequencies of 4-chloro-2-methylaniline in the temperature range 0-290 K are presented in Figure 3. The estimated frequencies at the extreme temperatures of all these compounds are given in Table 1. It can be seen that the estimated frequencies are in good agreement with the experimental values of  $\beta$ -phase in the case of 4-chloro-2-methylaniline. However the observed frequencies in 3-chloro-2-methylaniline having ortho- $\text{CH}_3$  substitution with respect to chlorine in the molecule differ by an extent of about one MHz from the estimated values.

## 4. Conclusion

4-Chloro-2-methylaniline is found to occur in two polymorphic phases namely  $\alpha$  and  $\beta$ .  $\alpha$ -Phase is a meta stable phase and is found to transform to a more stable  $\beta$ -phase with passage of time. NQR frequency estimation using Hammett's  $\sigma$ -values is not reliable in the case of ortho-substituted chloromethylanilines.

## References

- Biedenkapp D and Weiss A 1968 *J. Chem. Phys.* **49** 3933  
 Bray P J and Barnes R C 1957 *J. Chem. Phys.* **27** 551  
 Das T P and Hahn E L 1958 *Nuclear Quadrupole Resonance Spectroscopy* (New York: Academic Press)  
 McDonald A D and Hacopian S 1982 *J. Mol. Struct.* **83** 261  
 Setty D L S and Uchil J 1984 *Indian J. Pure Appl. Phys.* **22** 98