Crystal and molecular structure of dibenzofuran

ASOK BANERJEE

X-ray Laboratories, Presidency College, Calcutta-12

(Received 25 April 1972)

We are investigating the crystal and molecular structures of some heterocyclic compounds in our laboratory because of their widespread occurrence in nature as vitamins and pigments and their important use as drugs and dyes. Also, most of the cancer arresting substances are heterocyclic. Besides this, the diversity of reactions leading to heterocyclisation is an important problem of heterocycles and attempts have been made to correlate these with the structural information where available. In view of their importance, we have studied dibenzofuran, $C_{12}H_8O$. The preliminary report on this structure has already been reported by us in 1970. The present refined atomic parameters, bond angles and bond distances are consistent with similar other compounds. The molecule found to be planar has a plane of symmetry

X-RAY AND OPTICAL DATA

The colourless plate-like crystals of irregular shape were obtained by very slow evaporation from a solution in alcohol at room temperature ($\sim 27^\circ\text{C}$). Under polarizing microscope, the crystals show a perfect cleavage coinciding with the *ab* plane. The cleaved flake has birefringence of the order of 0.15, the two indices in the plane *ab* are 1.60 and 1.45 respectively. The cell constants were determined from oscillation and Weissenberg photographs (zero and upper level) about *a* and *b* axes with $\text{CuK}\alpha$ radiation. The density was measured by floatation method in ZnSO_4 solution. The crystal data are given in table 1.

Table 1

Compound	:	Dibenzofuran $C_{12}H_8O$
Unit cell	:	Orthorhombic
$a = 7.65\text{\AA}$	$b = 5.90\text{\AA}$	$c = 19.32\text{\AA}$
M.W. = 168.20	$d_{\text{calc.}} = 1.273 \text{ gm/cc}$	
MP = 87°C	$d_{\text{obs.}} = 1.265 \text{ gms/cc}$	
BP = 228°C	$V = 887\text{\AA}^3$	
$\mu = 6.39 \text{ cm}^{-1}$	$Z = 4$	
Space group : <i>Pnam</i>		
absent spectra ·		
$h0l, h = 2n+1$		
$0kl, k+1 = 2n+1$		

The integrated intensity of about 500 reflections were estimated visually from the zero and higher level equi-inclination Weissenberg photographs and put to same scale through cross layers. Lorentz and polarization corrections were applied but the correction for absorption was not done as the crystal was small and contained no heavy atoms ($\mu = 6.39 \text{ cm}^{-1}$). As the sample sublimes in air, a small crystal was sealed in a Lindemann glass capillary of 0.3 mm internal bore. The systematic absences of $h0l$ reflections when $h = 2n+1$ and $0kl$ reflections when $k+l = 2n+1$ give $Pnam$ and $Pna2_1$ as the possible space groups in this case. This ambiguity was solved by statistical method of Wilson. The distribution being centric, the space group is $Pnam$.

CRYSTAL STRUCTURE

The trial model was adopted by utilizing the existing knowledge of heterocyclic chemistry of similar compounds combined with refractive index and cleavage direction data. Approximate positional parameters were obtained by a combination of a weighted reciprocal lattice plot for $0kl$ reflections and the trial and error method. A least square refinement using the computer program ORFLS, written by Busing *et al* (1962), led to refinement of the structure with an R value of 0.145. Starting from an initial value of 40%, the final residual value obtained is 14.5% with 500 reflections collected along b -axis and scaled on a common base. The molecule is planar with the present state of refinement. A crystallographic mirror plane passes through the oxygen atom and is perpendicular to the molecular plane. The final values of the atomic parameters with the isotropic temperature factors for the non-hydrogen atoms are listed in table 2. Bond lengths and angles of the molecule are also shown in figure 1 to be consistent with existing accepted

Fig. 1

values. The distance of closest approach between carbon atoms in adjacent molecules is $3.77 \text{ \AA}$. This occurs between the atom C(1) of the molecule of x, y, z and the atom C(4) at $x, 1+y, z$. The distribution of the molecule inside the unit

cell explains the cleavage of the crystal very nicely. The residual value of 14.5% along with consistent values of bond lengths and angles justifies the correctness of the structure.

Table 2. Positional parameters, bond lengths, bond angles and temperature factors of non-hydrogen atoms

Atom	x/a	y/b	z/c	B
C(1)	0.2360	0.6992	0.3770	5.5
C(2)	0.1767	0.5456	0.4270	5.4
C(3)	0.0968	0.3404	0.4089	5.2
C(4)	0.0684	0.2803	0.3395	5.1
C(5)	0.2049	0.6334	0.3081	4.6
C(6)	0.1261	0.4299	0.2885	4.1
O(7)	0.2579	0.7655	0.2500	5.5

Table 3

Atoms	Bond lengths Å	Atoms	Bond angles
C(1)–C(2)	1.399	C(2)–C(1)–C(5)	114.8°
C(2)–C(3)	1.399	C(1)–C(2)–C(3)	122.0
C(3)–C(4)	1.404	C(1)–C(5)–C(6)	124.5
C(4)–C(6)	1.394	C(2)–C(3)–C(4)	121.5
C(1)–C(5)	1.407	C(3)–C(4)–C(6)	117.8
O(7)–C(5)	1.420	C(4)–C(6)–C(5)	119.4
C(6)–C(5)	1.395	C(6)–C(5)–O(7)	112.4
C(6)–C(6')	1.480	C(1)–C(5)–O(7)	123.1

A refinement of the structure considering anisotropic temperature factors and contributions of hydrogen atoms is in progress, and will be reported shortly, but the salient features of the crystal structure are not likely to change appreciably.

REFERENCES

- Banerjee A. 1971 *Proc. Seminars on Cryst., Centre of Advanced Study University of Madras*, B1, p. 8; 1971 *Proc. 58th Session Indian Sc. Congress. Part III No 42*, p 72; 1972 *Proc. Symposium Cryst.* Bhabha Atomic Research Centre, D1, p 18.
- Busing W. R., Martin K. O. & Levy H. A. 1962 *Publication ORNL-IM-305*, Oak Ridge National Laboratory, Tennessee.