

Rotational isomerism in methyl 2-chloro and methyl 2-bromopropionate

RANJIT DAS AND SAMIR KUMAR NANDY*

Optics Department,

Indian Association for the Cultivation of Science Jadavpur, Calcutta-700032

Infrared spectra of methyl 2-chloro and methyl 2-bromopropionate and the Raman spectra of the compounds at different temperatures in the liquid phase have been studied. Analysis of the results indicates that in the liquid state, molecules of each of the compounds exist in three stable rotameric forms viz, one *cis* and two *gauche*. The population of the *gauche* rotamers is observed to increase with lowering of temperature of the liquid.

1. INTRODUCTION

Molecules of some aliphatic esters and their derivatives have been shown to exist in the liquid phase in several configurations arising mainly from orientation about C—C bonds in them, the rotation about C—O bond being of minor importance. (Brown 1962, Neelkantan 1964, Charles *et al* 1973, Jones & Owen 1973). Identification of the rotamers of these compounds have been made mostly from partial analysis of their vibrational spectra (Josien & Calas 1955, Bellamy & Williams 1957, Brown 1962) and complete assignments of the observed frequencies have been reported only in a few cases. (Neelkantan 1964, Charles *et al* 1973). In order to provide adequate data needed for such assignments, a programme has been undertaken in this laboratory for studying the vibrational spectra of a series of halogenated esters. In the present communication, the results of investigation on the Raman and infrared spectra of two relatively simple haloesters viz, methyl 2-chloro and methyl 2-bromopropionate have been reported. Attempts have been made to make reasonable assignment of the observed spectra of the two molecules and to detect the existence, if any, of the conformers possible for each of the two molecules.

* Department of Physics, Jadavpur University, Calcutta-700032.

2. EXPERIMENTAL

The Raman and infrared spectra of the esters in the liquid state at ordinary temperature (90°C) were obtained photographically with a Fuess glass spectrograph and with a Perkin Elmer infrared spectrophotometer respectively. The Raman spectra of the liquids at low temperature (-110°C) were recorded using a bath of a mixture of alcohol and liquid nitrogen and that at high temperature (90°C) with a suitable electrical heater. The changes in the relative intensity of a chosen pair of Raman lines in each of the two liquids, with change of temperature, were measured by microphotometric methods.

3. RESULTS

The observed vibrational frequencies of the two compounds together with their relative intensities are listed separately in tables 1 and 2. These also contain the proposed assignments. Analysis of the results are presented in the following paragraphs.

4. DISCUSSION

The molecules of methyl 2-chloro and methyl 2-bromopropionate are structurally similar and if the OCH_3 group in each molecule is assumed to exist in the *cis*- configuration as in other esters (Neelkantan 1964, Charles *et al* 1973, Som and Mukherjee 1975, 1976, Som 1976) both of them will have only one rotational axis. Hindered rotation about this $\text{C}-\text{C}$ bond will give rise to three different conformers designated *cis*, *gauche 1* and *gauche 2* as shown in figure 1. The structural similarity of the two molecules is also reflected in the observed Raman and infrared spectra of the two compounds in the liquid phase.




Fig. 1. Three molecular conformations of methyl 2-chloro and methyl 2-bromo propionates

Table 1. Raman and infrared bands of methyl 2-chloropropionate in liquid state (30°C)


Raman shifts $\Delta\nu\text{cm}^{-1}$	Infrared bands νcm^{-1}	Assignments
134 (3)		
202 (1)		
243 (1) } 305 (4) }		δ (C - C - Cl)
356 (5)		δ (C - O - C)
404 (4)		δ (C - C - O)
465 (6)		δ (C - C - C)
623 (4)		π (C - O)
692 (8)	690 (m)	ν (C - Cl) } gauche
722 (6)		ν (C - Cl) }
771 (7)	765 (w)	ν (C - Cl)/CH ₂ Rock cis
859 (10)	860 (s)	(C - CH ₂) Rock
956 (8)	965 (m)	ν (C - C)
	985 (w)	ν (C - C)
997 (5)	1010 (s)	ν (C - C)
1077 (4)	1070 (s)	ν (-O - CH ₃)
	1100 (w) } 1165 (s) }	R(C - CH ₃)
1158 (4)	1195 (s)	(O - CH ₂) Rock
1240 (4)	1245 (s)	R(O - CH ₂)/(O - C )
1280 (5)	1285 (m)	CH wag
	1335 (m)	CH wag
	1385 (w)	δ (CH ₃) sym
	1440 (sh)	δ (OCH ₃) sym
1457 (8)	1450 (s)	δ (CH ₃) assym
1757 (8)	1745 (vs)	ν (C = O)
	2850 (m) } 2880 (w) }	ν (CH) in CH ₂ , sym
2879 (4)	2940 (sh) } 2960 (s) }	ν (CH) in CH ₂ , assym
2948 (8)		
2989 (7)	2990 (s)	ν (CH) in CH/ ν (CH) in OCH ₃

Table 2. Raman and infrared bands in methyl 2-bromopropionate in liquid state ,30°C)

Raman shifts $\Delta\nu\text{cm}^{-1}$	Infrared bands νcm^{-1}	Assignment
129 (5)		
239 (1) } 269 (9) } 330 (5)		δ (C-C-Br) δ (C-O-C)
370 (1)		δ (C-C-O)
442 (8)		δ (C'-C-C)
550 (6) (Thick)		π (C=O)/ ν (C-Br) } Gauche
653 (5)	654 (w)	ν (C-Br)
663 (8)	670 (m)	ν (C-Br) Cis
755 (6)	757 (m)	CH ₃ Rock
852 (8)	857 (s)	(C-CH ₃) Rock
956 (8)	963 (s)	ν (C-C)
995 (4)	1000 (s)	ν (C-C)
1065 (5)	1068 (s)	ν (O-CH ₃)
1094 (5)	1100 (m) }	(C-CH ₃) Rock
1158 (6)	1160 (s) }	R (C-CH ₃)
1190 (1)	1192 (w)	(O-CH ₃) Rock
1229 (6)	1226 (s)	R(O-CH ₃)/ ν (-O-C(=O)-)
	1272 (m)	CH wag
1304 (1)	1350 (s)	CH wag
1383 (2)		δ (CH ₃) sym
1443 (8)	1446 (vs)	δ (CH ₃) assym.
1748 (8)	1740 (vs)	δ (C=O)
2871 (5)		δ (CH) in CH ₃ sym.
2936 (6)	2957 (m) }	δ (CH) in CH ₃ assym.
2999 (6)	2985 (w) }	δ (CH) in CH/ ν (CH) in OCH ₃

A comparison of the data presented in tables 1 and 2 show that there are broad spectral regions in which the vibrational frequencies of the two molecules are similar. One of these regions extends from 3000 to 1300 cm^{-1} and another from 1300-900 cm^{-1} . From the known characteristics of the vibrational frequencies of the OCH_3 , CH_3 and CH groups, the bands at $\sim 3000 \text{ cm}^{-1}$ in each of the two molecules are attributed to the stretching vibration of the single CH bond and the Raman and infrared frequencies in the region 2960-2850 cm^{-1} are assigned to asymmetric and symmetric CH stretching vibrations in the methyl and methoxy groups.

The asymmetric CH deformation frequencies in these two groups are correlated with the frequencies at $\sim 1450 \text{ cm}^{-1}$ while the bands at $\sim 1380 \text{ cm}^{-1}$ are identified with the symmetric deformation mode. The *inplane* and *out of plane* wagging modes of the single CH give rise to the frequencies at ~ 1340 and 1280 cm^{-1} respectively. The strong Raman lines and infrared bands in the frequency region 1740-1757 cm^{-1} are easily assigned to the carbonyl stretching vibration. All these assignments are shown in the respective tables.

The region between 1250-900 cm^{-1} contains the frequencies arising from

$\text{O}-\text{CH}_3$ rocking and stretching, $\text{C}-\text{CH}_3$ rocking, $\begin{array}{c} \text{O} \\ || \\ -\text{C}-\text{O} \end{array}$ stretching and skeletal $\text{C}-\text{C}$ stretching modes of vibration. It is seen from the tables that the correspondence between these frequencies in the two molecules is very good. The bands at ~ 1200 and 1240 cm^{-1} are assigned to the $\text{O}-\text{CH}_3$ rocking modes while those at ~ 100 and 1160 cm^{-1} to similar modes of $\text{C}-\text{CH}_3$ group. However, the frequencies at $\sim 1240 \text{ cm}^{-1}$ may also be due to the $\text{C}-\text{O}$

$\begin{array}{c} \text{O} \\ || \\ -\text{O}-\text{C}- \end{array}$ stretching mode in the $\begin{array}{c} \text{O} \\ || \\ -\text{O}-\text{C}- \end{array}$ group. The vibration frequency at $\sim 1070 \text{ cm}^{-1}$ in both the compounds has been ascribed to another $\text{C}-\text{O}$ stretching mode in the $\text{O}-\text{CH}_3$ part of the esters. The Raman bands between 1000-950 cm^{-1} and their corresponding counterparts in the infrared spectra definitely represent the skeletal $\text{C}-\text{C}$ stretching frequencies.

Below the frequency range 900 cm^{-1} in molecules containing the $\text{C}-\text{CH}_3$ group, one or two bands representing $\text{C}-\text{CH}_3$ rocking mode have generally been observed and the bands 859 cm^{-1} in methyl 2-chloro and 852 cm^{-1} in methyl 2-bromopropionate are attributed to one such mode of vibration. The other mode is some what difficult to identify. In the chloro compound, there is a Raman line at 771 cm^{-1} (infrared band at 765 cm^{-1}) and in the bromocompound, a similar Raman line at 755 cm^{-1} (infrared) band at 757 cm^{-1}) is observed. The band in the former molecule may represent a frequency arising from the stretching of the $\text{C}-\text{Cl}$ bond, which is not possible in the latter. In 2-halobutanes, Benedetti and Cecchi (1972) have observed bands between 770-780 cm^{-1} which have been assigned by them to

a mode involving CH_2 rocking vibration. It, therefore, seems justified to correlate the frequencies 771 and 755 cm^{-1} in chloro and bromo esters respectively, with such vibrational mode. In the chloro compound, however, the superposition of this vibrational frequency with $\text{C}-\text{Cl}$ stretching frequency is highly probably.

The frequency region from about 600 cm^{-1} down to $\sim 100\text{ cm}^{-1}$ has been attributed mainly to vibrational modes arising from the deformation of the molecular skeleton. Though some degree of correspondence in the spectra of the two compounds is found also in this respect, the correlation is not very certain. Following the assignments made in many molecules containing similar skeleton (Wilmshurst 1957, Charles *et al* 1973) some of the observed vibration frequencies have been identified and are shown in the tables without further elaboration but some discussion regarding the $\text{C}=\text{O}$ deformation vibration are presented.

The $\text{C}=\text{O}$ out of plane bending frequencies, designated as $\pi(\text{C}=\text{O})$, in ester molecules lie usually in the region $625\text{-}500\text{ cm}^{-1}$. In methyl acetate (Wilmshurst 1957), the $\pi(\text{C}=\text{O})$ frequency is reported at $\sim 615\text{ cm}^{-1}$ and according to Thompson (1945) this frequency in the alkyl propionates occurs at $\sim 600\text{ cm}^{-1}$. Som (1975) assigned the frequency 617 cm^{-1} in ethyl 2-chloropropionate and Som and Mukherjee (1976) attributed that at 536 cm^{-1} in methyl 3-bromopropionate to this mode of vibration. In the Raman spectrum of methyl 2-chloropropionate, there is a band at 623 cm^{-1} and in methyl 2-bromopropionate a broad Raman line at 550 cm^{-1} has been observed.

The former is easily identified with the $\pi(\text{C}=\text{O})$ mode of vibration while the latter is interpreted as arising from superposition of the frequency of the $\pi(\text{C}=\text{O})$ vibration and a $\text{C}-\text{Br}$ stretching vibrational frequency. This is similar to that observed in methyl 3-bromopropionate. (Som & Mukherjee).

A careful examination of the data given in tables 1 and 2, show that the total number of bands observed in these spectral regions, whose assignment has been presented in the above paragraphs, is barely enough to account for all the vibrational modes possible for only one molecular configuration in each case. But the spectral region between $800\text{-}650\text{ cm}^{-1}$ in methyl 2-chloropropionate and $700\text{-}550\text{ cm}^{-1}$ in methyl 2-bromopropionate contain a number of extra bands which points to the existence of rotational isomerism in these molecules. It can be seen from table 1 that in the Raman spectrum of methyl 2-chloropropionate, there are three strong lines at 771 , 722 and 692 cm^{-1} which occur in the region specified for the carbon-chlorine stretching frequencies in alkyl chlorides (Shipman *et al* 1962) and in some chloro-substituted esters (Neelkantan 1964, Som 1976). In view of this, all the three frequencies have been attributed to $\text{C}-\text{Cl}$ stretching vibrations. Similarly,

following the assignments given in the cases of the alkyl bromides (Bentley), ethyl bromoacetate (McBee and Cristian 1955) and methyl 3-bromopropionate (Som and Mukherjee 1976), the three frequencies 663, 653 and 550 cm^{-1} observed in the Raman spectrum of methyl 2-bromopropionate are considered as arising from carbon-bromine stretching vibrations. The Raman line at 771 cm^{-1} in the chlorocompound and 550 cm^{-1} in the bromocompound, may, as shown earlier, also be due to CH_2 rocking mode and π ($\text{C} = \text{O}$) mode of vibration of the respective molecules.

The analysis of the special region representing frequencies due to carbon-halogen stretching vibration thus reveals that there are three such frequencies in each of the two molecules and this leads to the conclusion that the molecules of the two compounds, in the liquid state, exist in at least three different conformations, earlier termed *cis*, *gauche* 1 and *gauche* 2. It has been found that in many haloacetyl halides (Khan and Jonathan 1969) the $\text{C}-\text{Cl}$ stretching frequency for the *cis* isomer (*trans* w.r.t to halogen) occur at a value higher than that due to the *gauche* isomer and the same trend is observed also in the case of chloroacetone (Mizushima 1954), ethyl haloacetates (Neelkantan 1964) and ethyl halopropionate (Som 1976). In view of this, of the three carbon-halogen stretching frequencies in methyl 2-chloro and methyl 2-bromopropionate, the one of highest value, in each case, has been attributed to the *cis* form and the remaining two frequencies to the two *gauche* forms of the molecules.

In order to study the relative stability of the rotamers of each of the compounds, measurements on the temperature variation of the relative intensities of a pair of Raman lines due to carbon-halogen stretching in two different structural forms have been made over the temperature range of 90°C to -110°C. In the case of methyl 2-chloropropionate molecule, it has been observed that as the temperature decreases, the intensity of Raman line at 771 cm^{-1} due to the *cis* form decreases while of the Raman line 722 cm^{-1} due to one of the two *gauche* forms increases. Similarly, in methyl 2-bromopropionate, the ratio of the intensities of the Raman line 663 cm^{-1} of the *cis* rotamer to that of 653 cm^{-1} representing one of the *gauche* rotamers has been found to decrease with decrease of temperature. These facts lead to the conclusion that in each of the methyl 2-halopropionates, in the liquid state, the population of one of the two *gauche* rotamers, increases with lowering of temperature or in other words, in the liquid phase of each of the two compounds, this *gauche* rotamer is more stable than the *cis* rotamer. The conformation of the molecules in the solid state could not be ascertained because perfect crystalline masses were not obtained in spite of repeated attempts. However, it may be of interest to note that in the similar molecule of ethyl 2-chloropropionate, the *gauche* rotamer has been found to be the only conformer present in the crystalline phase at low temperature (Som 1976).

ACKNOWLEDGEMENT

The authors are grateful to Prof. G. S. Kastha, D.Sc. for his kind guidance in the progress of the present work.

REFERENCES

- Bellamy I. J. & Williams R. L. 1957 *J. Chem. Soc.* 4294.
Benedetti E & Cecchi P 1972 *Spectrochim Acta.* **28A**, 1007
Bentley F. F. McDevitt N. T & Rozeck A. L. 1964 *Spectrochim Acta.* **20**, 105.
Brown T. L. 1962 *Spectrochim Acta.* **18**, 1615
Charles S. W. Jones G. I. L & Owen N. L. 1973 *J. Chem. Soc. Part II((Trans. Farad)* **69**, 1454.
Jones G. I. L & Owen N. L. 1973 *J. Mol. Struct* , **18**, 1.
Jones M. L. & Calas R. 1955 *Compt Rend.* **240**, 1641.
Khan A & Jonathan N. 1969 *J Chem Phys.*, **50**, 1801
McBee E T & Cristian D. L. 1955 *J Am. Chem. Soc* **77**, 755.
Mizushima S. 1954 *Structure of Molecules and Internal Rotation*, Academic Press Inc. New York., 73.
Neelkantan P. 1964 *Proc Indian Acad. Sci., Sec A.*, **59**, 385.
Shipman J. J., Folt V. L. & Krimm S., 1962 *Spectrochim Acta* **18**, 1603.
Som J & Mukherjee D. K. 1975 *Indian J. Pure Appl. Phys.*, **13**, 348.
Som J & Mukherjee D. K. 1976 *Indian J Pure Appl Phys.* **14**, 151.
Som J. 1976 *Indian J. Pure. Appl. Phys.*, **14**, 313.
Thompson H. W & Torkington P 1945 *J. Chem. Soc* , 640.
Wilmschurst J. K. 1957 *J. Mol. Spectry* , **1**, 201.