Hypersensitive transitions in Nd^{3+} complexes

P. P. VAISHNAVA AND S. P. TANDON

Department of Physics, University of Jodhpur, Jodhpur

The transition ${}^4\text{G}_{5/2} \rightarrow {}^2\text{G}_{7/2} \leftarrow {}^4\text{I}_{9/2}$ obeying the selection rule $\Delta J \leq 2$, known as hypersensitive transition, has been studied for some Nd^{3+} complexes. Various theories put forward to explain the phenomenon have been discussed. The results indicate that the hypersensitivity may be explained by means of vibronic mechanism with the inclusion of covalency.

1. INTRODUCTION

The absorption spectra of rare earth complexes have been a subject of several investigation¹⁻⁵. In Nd^{3+} complexes, the transition $({}^4\text{G}_{5/2}, {}^2\text{G}_{7/2}) \leftarrow {}^4\text{I}_{9/2}$ obeying the selection rule $\Delta J \leq 2$ is known as hypersensitive transition. These bands give rise to intense bands⁶. In these transitions the intensity of corresponding bands is changed by changing the ligand or solvent or both. The explanation of the phenomenon of hypersensitivity has proved to be an intriguing problem. The present communication deals with the various theories advanced to explain this phenomenon in the light of recent studies of spectral intensities.

2. CALCULATION AND RESULTS

The oscillator strength, P , for intra- f^n transitions of rare earths is given by Judd⁷ as

$$P = \sum_{\lambda=2,4,6} \tau_{\lambda} \nu (f^n \Psi_J || U(\lambda) || f^n \Psi'_{J'})^2 / (2J+1), \quad \dots \dots (1)$$

where ν is the energy (in cm^{-1}) of the transition $\Psi_J \rightarrow \Psi'_{J'}$ and $U(\lambda)$ is a tensor operator of rank λ . The three quantities τ_{λ} may be treated as parameters. The experimental oscillator strength, P_{exptl} values have been found⁷ to be due to the contributions from induced electric-dipole (P_{ed}), magnetic-dipole (P_{md}) and electric-quadrupole (P_{eq}) modes. Since P_{eq} ($\sim 10^{-11}$) and P_{md} ($\sim 10^{-8}$), are negligible⁶ in comparison to P_{exptl} ($\sim 10^{-6}$), they may be neglected. Hence P_{exptl} values may be substituted in place of P in eq. (1).

It has been found⁸ that in general, the oscillator strengths for Nd^{3+} complexes can be expressed in terms of τ_1 and τ_6 parameters only. But it is important to note that the effect of solvent or ligand on the intensity of a hypersensitive transition can be explained by τ_2 parameter only. The values of τ_2 parameter have been given in Table 1.

The explanation of the phenomenon of hypersensitivity has proved to be an intriguing experimental and theoretical problem. Karriaker^{9,10} has studied the hypersensitive transitions in six-, seven- and eight-coordinated Nd^{3+} , Ho^{3+} and Nd^{3+} β -diketonates and has shown that the changes in the oscillator strengths of these transitions may be correlated with the changes in the coordination and symmetry of the rare earth ion. Menzel and Gruber¹¹ have shown that the external electric field may be useful tool in the study of hypersensitive transitions. The following proposals have been advanced for the explanation of these transitions:

(a) *Pseudo-quadrupole* transitions:

Jorgensen and Judd⁶ have called these transitions as Pseudoquadrupole transitions and showed their existence due to an asymmetric distribution of dipoles induced by electromagnetic radiation in the medium surrounding the rare earth ions. These dipoles produce a large variation in the electric vector across the ion which results in the enhancement of the electric-quadrupole transition. However, their results are based upon many approximations.

(b) Inclusion of spherical harmonics in the crystal field potential:

This mechanism, advanced by Judd⁷, results from the absence of a center of inversion in the compound. In symmetries where a center of inversion is absent, only transitions obeying the selection rule $\Delta J \leq 2$ are affected by the spherical harmonics $Y_1^{(m)}$ which mix the $4f^n$ and $4f^{n-1} 5d^1$ configurations. Thus, according to this model the presence of harmonics $Y_1^{(m)}$ is essential in the crystal field of the complex. In NdA_3 the phenomenon of hypersensitivity is observed even though $Y_1^{(m)}$ is not permitted. Hence this model is not satisfactory.

(c) Polarization of the surrounding ions:

Blasse and Brill¹² have shown that hypersensitivity of the ${}^7F_2 \leftarrow {}^5D_0$ transition of the emission of Eu^{3+} in garnets can be correlated with the state of polarization of the surrounding anions. The polarizing field at the anions can be taken to be equal to the difference, Δ , between the average ionic charges on garnet structure (viz., tetrahedral and octahedral) sites. It has been found¹² that as Δ increases, the relative intensity of hypersensitive ${}^7F_2 \leftarrow {}^5D_0$ transitions increases, whereas the intensity of ${}^7F_1 \leftarrow {}^5D_0$ transition remains constant.

(d) Double-Photon transitions :

According to Axe¹³ the *double-photon* transitions may be the possible cause of hypersensitivity. In this process the spectral intensity of the transition should be dependent on the intensity of the incident radiations. To test this the absorption of Nd³⁺ p-nitrobenzoic acid in the spectral region 5000-6500 Å as a function of the intensity of incident radiation was studied. The results showed no change in the values of the oscillator strengths of the hypersensitive transitions. Hence, the *double-photon* transitions mechanism cannot explain the cause of hypersensitivity.

(e) Vibronic mechanism with the inclusion of covalency :

Recently, Henrie and Choppin¹⁴ have proposed that the phenomenon of hypersensitivity may be explained by the inclusion of covalency within the framework of vibronic mechanism. According to this mechanism the oscillator strength of the hypersensitive transitions should vary approximately as νb^2 , where the degree of covalency ($b^{1/2}$) has been defined by ¹⁴

$$b^{1/2} = \left(\frac{1-\beta}{2} \right)^{1/2}, \quad \dots\dots\dots (2)$$

in which β is known as nephelauxetic ratio and is given by¹¹

$$\beta = \frac{\nu_{\text{comp}}}{\nu_{\text{aquo}}}, \quad \dots\dots\dots (3)$$

where ν_{comp} and ν_{aquo} are the energies of the hypersensitive transitions of Nd³⁺ complex and aquo ion respectively.

Table 1. Spectral data for Nd³⁺ complexes

Nd ³⁺ Complexes	ν (cm ⁻¹)	P_{exptl} $\times 10^6$	$\tau_2 \times 10^8$ (cm)	b^2 $\times 10^2$
Aquo ion	17300	8.00	0.30	—
NBA	17290	26.22	9.70	1.7
A ^a	17167	33.39	2.18	6.2
TFA ^a	17167	44.90	2.85	6.2
BA ^a	17105	53.02	3.37	7.5

The computed values of $b^{1/2}$ together with the energies and P_{exptl} values of the hypersensitive transitions have also been given in Table 1. A plot of

P vs νb^2 for (${}^4G_{5/2}$, ${}^2G_{7/2}$) $\leftarrow {}^4I_{9/2}$ transitions of Nd^{3+} complexes (Fig. 1) indicates approximately linear relationship. Thus, our results seem to indicate,

Fig. 1.

in agreement with those of Henri and Coppin, that the phenomenon of hypersensitivity may be explained by means of a vibronic mechanism with the inclusion of covalency.

ACKNOWLEDGEMENT

Authors express their thanks to the Incharge Computer Centre, University of Jodhpur for providing computer facilities and to Dr. A. N. Nigam for his keen interest and the encouragement.

REFERENCES

1. G. H. Dicke, *Spectra and Energy levels of Rare Earth ions in Crystals* (John Wiley & Sons, Inc., New York, 1968).
2. S. P. Tandon and P. C. Mehta *J. Chem. Phys.* **52**, 4896, 5417 (1970).
3. S. P. Tandon, P. P. Vaishnava and R. C. Mathur *Sol. State Comm.* **15**, 787 (1974).
4. P. P. Vaishnava, S. P. Tandon and M. P. Bhutia, *Spectry Letters* **7**, 515 (1974).
5. M. P. Bhutia, P. P. Vaishnava and B. S. Sankhla *Can. J. Spectry.* **20**, 77 (1975).
6. C. K. Jorgensen and B. R. Judd, *J. Mol. Spectry.* **8**, 281 (1964).
7. B. R. Judd, *J. Chem. Phys.* **44**, 839 (1966).
8. D. G. Karrakar *Inorg. Chem.* **6**, 1863 (1967).
9. D. G. Karrakar *Inorg. Chem.* **7**, 473 (1968).
10. E. R. Menzel and J. B. Gruber *J. Chem. Phys.* **52**, 1008 (1970).
11. G. Blasse and A. Bril *J. Chem. Phys.* **47**, 5442 (1967).
12. J. D. Axe Jr., *Phys. Rev.* **136**, A. 42 (1964).
13. D. E. Henrie and G. R. Choppin *J. Chem. Phys.* **49**, 47 (1968).