

Crystal and molecular structure of 2-phenyl-3N-[(3''-chloro-4''-fluoro)phenyl]-4-thiazolidinone

K V Arjuna Gowda, M K Kokila and Puttaraja

Department of Physics, Jnanabharathi Campus, Bangalore University,
Bangalore-560 056, India

M V Kulkarni

Department of Chemistry, Karnataka University, Dharwad-580 003, India

and

N C Shivaprakash*

Department of Instrumentation Indian Institute of Science,
Bangalore 560 012, India

E-mail : shiv@isu.isc.ernet.in

Received 25 July 2001, accepted 10 August 2001

Abstract The crystal and molecular structure of 2-phenyl 3N-[(3''-chloro 4''-fluoro) phenyl] 4 thiazolidinone have been determined by X ray structure analysis. They belong to the family of thiazolidinones. The structure was solved by direct method. The molecular packing of the non planar molecules are held by Van der Waals interactions.

Keywords Thiazolidinone, X-ray crystallography

PACS Nos. 61.66.Fn, 61.10.-i

1. Introduction

The title compound belongs to the family of thiazolidinones. 2-aryl-1, 3-thiazolidin-4-ones have been found to exhibit anti-fungal, anti-tubercular, anti-viral and muscle relaxant properties [1]. 3-(2-pyridyl)-2-(2-tolyl) thiazolidin-4-one was found to exhibit anti-bacterial property [2] and subsequently its crystal structure was reported [3]. 2-(3-pyridyl) thiazolidin-4-ones with different substituents at position 5 have emerged as a new class of platelet factor antagonists [4]. The compound under study was found to show considerable inhibition (33%) of the carrageenan induced edema in rat paw [5]. It was reported by our group that 3-chloro-4-fluoro benzylidene aniline with p-chloro group in the benzylidene ring adopts a twisted conformation [6]. It has been recently pointed out that organo fluorine has low proton affinity and hardly accepts hydrogen bonds [7].

Structural investigation of the title compound has been taken up in view of the fact that it contains three ring systems and it is

of interest to study the arrangement of these which the physical method (IR-PMR) do not provide. Further, it is also worth considering the changes in the conformation across C-N bond when the two phenyl rings are intervened by a thiazolidinone ring.

2. Experimental

Crystals suitable for X-ray diffraction were grown by slow evaporation technique using Ethanol. X-ray intensity data was collected on CAD4 diffractometer with $\text{CuK}\alpha$ radiation. Three dimensional intensity data were corrected for Lorentz and Polarisation effects. Absorption and extinction corrections were not applied. The crystallographic data and other details of experimental and refinement are summarized in Table 1.

The structure was solved by direct method using SHELXS-97 [8] program. The structure thus obtained was refined using SHELXL-97 [9] program by a full-matrix least-squares method.

* Corresponding Author

A sigma weighting scheme was applied and the refinement was continued till final value of $R = 0.041$, $R_w = 0.117$.

Table 1. Crystal data for the title compound

Crystal morphology	Colourless needle like
Crystal size	0.2 × 0.2 × 0.15 mm
Chemical formula	C ₁₅ H ₁₁ NOSClF
Molecular weight	307.76
Crystal system	Trichinic
Space group	P $\bar{1}$
Cell constants	$a = 6.019(1)$ Å $b = 10.256(1)$ Å $c = 11.549(3)$ Å $\alpha = 75.50(1)^\circ$ $\beta = 78.47(2)^\circ$ $\gamma = 76.08(1)^\circ$
Volume	662.50 (2) Å ³
Number of formula units Z	2
Density (calculated) D_c	1.543 gm/cm ³
Unique data with measured	2516
Observed data with $I \geq 4\sigma(I)$	2423
$F(000)$	316
R	0.041

3. Results and discussion

The fractional coordinates and equivalent isotropic temperature factors for non-hydrogen atoms are presented in Table 2, bond

Table 2. Fractional coordinates and isotropic temperature factors U_{iso} (Å²) for non-hydrogen atoms with e.s.d's are in the parentheses

Atom	U_{iso}			
Cl	0.5097(1)	0.2168(1)	0.4901(1)	0.0498(2)
S1	0.1028(1)	0.8290(1)	-0.0099(1)	0.0374(2)
F	0.8871(3)	0.3520(2)	0.4891(2)	0.0565(5)
O1	0.2843(3)	0.4384(1)	0.0906(2)	0.0421(5)
N3	0.3277(3)	0.6323(2)	0.1420(2)	0.0276(4)
C1''	0.4703(4)	0.5612(2)	0.2316(2)	0.0268(5)
C2''	0.4227(4)	0.4394(2)	0.3094(2)	0.0308(5)
C3''	0.5635(4)	0.3689(3)	0.3951(2)	0.0335(5)
C4''	0.7499(4)	0.4203(3)	0.4049(2)	0.0365(5)
C5''	0.7968(4)	0.5406(3)	0.3307(2)	0.0376(6)
C6''	0.6599(4)	0.6104(3)	0.2427(2)	0.0324(5)
C2	0.3254(4)	0.7794(2)	0.0892(2)	0.0279(5)
C1'	0.2668(4)	0.8735(2)	0.1767(2)	0.0277(5)
C2'	0.4099(5)	0.9638(3)	0.1694(2)	0.0370(6)
C3'	0.3488(6)	1.0609(3)	0.2406(3)	0.0490(7)
C4'	0.1449(6)	1.0673(3)	0.3201(3)	0.0499(7)
C5'	0.0036(5)	0.9760(3)	0.3298(2)	0.0450(7)
C6'	0.0627(4)	0.8798(3)	0.2587(2)	0.0350(5)
C5	0.1535(4)	0.6582(3)	-0.0341(2)	0.0376(6)
C4	0.2580(4)	0.5632(2)	0.0716(2)	0.0309(5)

lengths and bond angles in Table 3 respectively. The general view of the molecule indicating the numbering scheme is as shown in Figure 1. An ORTEP [10] plot of the molecule with probability thermal ellipsoids is shown in Figure 2.

Table 3. Selected bond distances (Å) and bond Angles (°)

Atom	Bond length	Atom	Angle
Cl-C3''	1.728(3)	C5-S1-C2	91.6(1)
S1-C5	1.787(3)	C4-N3-C1''	-121.3(2)
S1-C2	1.832(2)	C4-N3-C2	115.0(2)
F-C4''	1.349(3)	C6''-C1''-C2''	119.2(2)
O1-C4	1.219(3)	C2''-C3''-C1	120.6(2)
N3-C4	1.381(3)	C4''-C3''-C1	119.7(2)
N3-C1''	1.420(3)	C6''-C1''-N3	120.8(2)
N3-C2	1.476(3)	C2''-C1''-N3	120.0(2)
C1''-C6''	1.392(3)	C3''-C2''-C1''	120.1(2)
C1''-C2''	1.399(3)	C2''-C3''-C4''	119.7(2)
C2''-C3''	1.379(3)	F-C4''-C5''	119.2(2)
C3''-C4''	1.384(4)	F-C4''-C3''	119.8(2)
C4''-C5''	1.370(4)	C5''-C4''-C3''	120.9(2)
C5''-C6''	1.383(4)	C4''-C5''-C6''	119.8(2)
C2-C1'	1.503(3)	C5''-C6''-C1''	120.3(2)
C1'-C2'	1.386(3)	N3-C2-C1'	116.4(2)
C1'-C6'	1.392(3)	N3-C2-S1	103.9(1)
C2'-C3'	1.384(4)	C1''-N3-C2	119.5(2)
C3'-C4'	1.376(4)	C1'-C2-S1	108.7(1)
C4'-C5'	1.381(4)	C2'-C1'-C6'	119.0(2)
C5'-C6'	1.376(4)	C2'-C1'-C2	119.3(2)
C5-C4	1.501(4)	C6'-C1'-C2	121.4(2)
		C3'-C2'-C1'	120.6(2)
		C4'-C3'-C2'	119.8(3)
		C3'-C4'-C5'	120.0(2)
		C6'-C5'-C4'	120.5(4)
		C5'-C6'-C1'	120.1(2)
		C4-C5-S1	107.2(1)
		O1-C4-N3	124.0(2)
		O1-C4-C5	123.3(2)
		N3-C4-C5	112.6(2)

The carbonyl oxygen attached to C4 is almost eclipsed with a dihedral angle of $10.3(3)^\circ$. The two phenyl rings are cis to C2-N3 bond and are in gauche arrangement with C1''-N3-C2-C1' (torsional angle of $55.4(3)^\circ$) and the thiazolidinone ring is non-planar (C2 and C5 are deviated from the plane of the thiazolidinone ring by 0.261 (2) and 0.259(3) Å respectively, torsional angle C5-S1-C2-N3 of $-28.3(2)^\circ$).

The C-C and C-H bond distances in the two phenyl rings are in the normal range of 1.37 to 1.39 Å and 0.92 to 0.93 Å respectively. The N3-C4 bond length 1.381(3) Å is shorter than

N3-C1'' bond length 1.420(3)Å due to two delocalisation of the nitrogen lone pair on the carbonyl group. The C5-H5 and C2-H2 bond distances are typical of Csp³ values and are found to be 0.97 to 0.98Å respectively. The C3''-C1 bond length 1.728(3)Å




Figure 1. Perspective view of the molecule with atomic numbering scheme

and C4''-F bond distance 1.349(3) Å are similar to those observed in benzylidene anilines [6]. The C2-S1 bond distance (1.832(2) Å) is close to and C5-S1 bond distance (1.787(3)Å) is less than the C-S single bond distance (1.832Å) [11] but still indicates single bond character [12]. The C4-O1 bond distance (1.219(3)Å) is almost comparable with C=O distance (1.214(2)Å) [13].




Figure 2. ORTEP plot of the molecule viewed along *b* axis

4. Molecular packing

The packing diagram of the molecule viewed along *a* axis is shown in Figure 3. The molecule on the whole is non-planar. The orientation of the benzene ring by 38.9(1) and that of phenyl ring by 84.9(1)° may be due to the presence of thiazolidinone ring. The molecules are packed in parallel layers in *bc* plane. The

hydrogen bond and a few H-atom interactions are listed in Table 4.




Figure 3. Packing diagram of the molecule viewed along

Table 4. Hydrogen bond and a few H atom interactions

D	H	A	D-H	H A	D A	D-H A	Symm
C2''	H2''	O1	0.930	2.371	2.817(3)	106.47	1_555
C2''	H2''	F	0.930	2.296	3.645(3)	129.28	1_555
C5''	H5''	F	0.930	2.694	3.594(4)	142.41	2_344
C4'	H4'	F	0.930	2.705	3.767(3)	168.21	1_465
C5	H5A	O1	0.970	2.198	3.257(4)	166.65	2_545
C5	H5B	O1	0.970	2.672	3.271(3)	116.64	2_445

Symm (1) *x*, *y*, *z*; (2) *-x*, *-y*, *-z*

Acknowledgment

One of the authors Shri. K. V Arjuna Gowda gratefully thank Dr L. V. G. Nargund and his student M. Srinivasmurthy, Dr. Nargund Research Foundation, Bangalore for supplying powder form sample.

References

- [1] G R Newkome and A Nayak *Advances in Heterocyclic Chemistry* Vol. 25 (eds) A R Katritzky and A J Boulton p 83 (New York Academic) (1979)
- [2] G Fenech *Farmaco* Ed. Sci 17 556 (1962)
- [3] D Hickel, J M Leger, A Carpy, M G Vigorita, A Chimri and S Grasso *Acta Cryst* C39 240 (1983)
- [4] Y Tanabe, H Yamamoto, M Murakami, K Yanagi, Y Kubota, H Okumura, Y Sanemitsu and G Suzukamo *J Chem Soc Perkin, Trans I* 935 (1965)

- [5] L V G Nargund and M Srinivasmurthy *M. Pharm Dissertation* (Al-Ameen College of Pharmacy, Bangalore University, Bangalore, India) (1994)
- [6] K V Arjuna Gowda, V Prakash, M K Kokila, Puttaraja, M V Kulkarni and N C Shivaprakash *Z. Krist.* **214**, 857 (1999)
- [7] T N Guru Row *Co-ordination Chemistry Reviews* **183** 81 (1999)
- [8] G M Sheldrick *SHELXS97 Program for the Solution of Crystal Structures* (University of Gottingen, Germany) (1997)
- [9] G M Sheldrick *SHELXL97 Program for the Refinement of Crystal Structures* (University of Gottingen, Germany) (1997)
- [10] C K Johnson *ORTEP REPORT ORNL-3794* (Oak Ridge National Laboratory, Tennessee) (1965)
- [11] L E Sutton *Interatomic Distances and Configurations in Molecules and Ions* (Supplement) (London: The Chemical Society) (1995)
- [12] M K Kokila, Puttaraja, M V Kulkarni and Sarala Thampi *Acta Cryst.* **C51** 330 (1995)
- [13] T N Chandrakantha, Puttaraja, M K Kokila and N C Shivaprakash *Acta Cryst.* **C54** 1685 (1998)