

The prediction of mass of Z' -boson in an $SO(10)$ -based model

S Sahoo

Department of Physics, National Institute of Technology (Deemed University)
Durgapur-713 209, West Bengal, India

E-mail: sskadevsahoo@yahoo.com

Received 29 May 2005, accepted 22 November, 2005

Abstract Z' -bosons are known to naturally exist in well-motivated extensions of the Standard Model. The mass of Z' -boson is considered in an $SO(10)$ -based model. We have used the low energy parameters and the estimated value so obtained is accessible to experiments at low energies.

Keywords Z -boson, Z' -boson, Higgs boson, grand unified theory

ACS Nos. 14.70 Hp; 14.70 Pw; 14.80.Bn; 12.10 Dm

1. Introduction

The presence of additional Z' -boson is predicted in a certain class of Grand Unified Theories (GUTs) [1] with a gauge group whose rank is higher than that of the Standard Model (SM). In additional GUTs, the scale of the Z' mass is arbitrary. However, in perturbative heterotic string models with supergravity mediated supersymmetry breaking, the $U(1)'$ and electroweak breaking are both driven by a radiative mechanism, with their scales set by the soft supersymmetry breaking parameters, implying that the Z' mass should be less than around 1 TeV [3]. The additional $U(1)'$ gauge boson Z' can mix with the hypercharge $U(1)_Y$ gauge boson through the kinetic term at above the electroweak scale, and also it can mix with the Standard Model Z -boson after the electroweak symmetry is spontaneously broken. Through those mixings, the Z' -boson can affect the electroweak observables at the Z -pole and W -boson mass m_W . Both Z - Z' mixing and Z' contribution can be explored directly in experiment [4]. There has also been the study on the effect of Z' -boson in the electroweak phase transition [5] with the mass of Z' -boson $m_{Z'} = 400$ GeV. There is a strong first-order electroweak phase transition with reasonable large value of Higgs mass to explain cosmological phenomena of Baryogenesis in the electroweak scale rather than GUT scale. However, there are stringent limits on the mass of an extra Z' from the non-observation of direct production followed by decays into e^+e^- or $\mu^+\mu^-$ by the Collider Detector at Fermilab (CDF) [6], while

indirect constraints from the precision data also limit the Z' mass (weak neutral current processes and Large Electron Positron Collider (LEP II)) and the Z - Z' mixing angle θ severely [7,8]. These limits are model-dependent, but are typically in the range $m_{Z'} > O(500)$ GeV and $|\theta| < few \times 10^{-3}$ for standard GUT models. A Z' could be relevant to the NuTeV experiment [9] and, if the couplings are not family universal [7,10], to the anomalous value of the forward-backward asymmetry A_{FB}^b [8]. (Earlier hints of a discrepancy in atomic parity violation have largely disappeared due to improved calculations of radiative corrections [11].) There is, thus, both theoretical and experimental motivation for an additional Z' , most likely in the range of 500 GeV - 1 TeV. Also, in this mass range, it should be possible to carry out significant diagnostic probes of Z' couplings at the Large Hadron Collider (LHC) and at a future Next Linear Collider (NLC) [3], which would complement those from the precision experiments [8]. The existence of the Z' -boson would also suggest a spectrum of particles considerably different than most versions of the Minimal Supersymmetric Standard Model (MSSM) [12]. Here, we investigate the mass of Z' -boson in $SU(2) \times U(1) \times U(1)$ model, which is a subgroup of $SO(10)$.

This paper is organized as follows: in Section 2, we give a brief account of the $SU(2) \times U(1) \times U(1)$ model, which is a subgroup of $SO(10)$. When the symmetry group breaks down to $U(1)_{em}$, the additional Z' -gauge boson acquires mass. We also

discuss the mass relations between particles in the model. In Section 3, we evaluate the mass of Z' -boson using the experimental values of parameters. We summarize the result at the end.

2. The model and mass relations

The Standard Model of the electroweak and strong interactions describes nearly all-experimental data available today [13]. The SM is based on the gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$, or G_{321} . In 1974, Georgi and Glashow [1] showed that the smallest simple gauge group G , which can contain the SM gauge group, is $G = SU(5)$. The number n of neutral gauge bosons of a GUT is given by $n = \text{rank}[G]$. We have $\text{rank}[SU(5)] = 4$. Therefore, there is no room for additional neutral gauge bosons in the $SU(5)$ GUT. All GUTs with gauge groups larger than $SU(5)$ predict atleast one extra neutral gauge boson Z' . It was shown by Fritzsch and Minkowski in 1975 [1] that the next interesting gauge group larger than $SU(5)$ is $SO(10)$. The group $SO(10)$ has rank 5, so the $SO(10)$ GUT does predict the existence of a Z' . Also, since $SO(10)$ has a larger rank than that of G_{321} , there are several ways in which it can be broken down to G_{321} . For example,

$$SO(10) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X, \quad (1)$$

$$\rightarrow SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1). \quad (2)$$

The first chain leads to the additional Z' -gauge boson [14], while the second chain yields the Left-Right Symmetric Model (LRSM) [15] which extends the SM gauge group to $SU(2)_L \times SU(2)_R \times U(1)$ with a right-handed charged boson as well as an additional neutral current.

Let us consider eq. (1), which is our desired model. While the symmetry group breaks down to $SU(3)_C \times U(1)_{em}$, the additional Z' gauge boson acquires mass. In the symmetry breaking scheme of eq. (1), the Z' is conventionally called Z_χ . The generator of the Z_χ commutes with the $SU(5)$ generators, so the fermionic couplings of the Z_χ are the same for all fermions in an $SU(5)$ multiplet. The mass of the Z_χ depends on the structure of the Higgs sector: m_{Z_χ} may be up to $O(m_{GUT})$. For at least one choice of Higgs bosons [14], the Z_χ may have a mass as low as a few hundred GeV. Our calculations will also prove this statement later in Section 3.

The Higgs doublet of the Standard Model is given by [16]

$$\Phi = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi + i\phi_3 \end{pmatrix}, \quad (3)$$

where ϕ_1, ϕ_2 , and ϕ_3 are the Goldstone bosons. Let us consider the field ϕ has a non-zero vacuum expectation value, $\langle \phi \rangle = k$.

The vacuum expectation value k is directly related to the effective strength of low-energy weak interactions [14]. Experimentally, it is shown that $k = 246$ GeV. The symmetry breaking $SU(2)_L \times U(1)_Y \rightarrow U(1)_{em}$ takes place because of the non-zero value of the vacuum expectation value of the field $k = 246$ GeV [17]. This vacuum expectation value can be determined by the minimization of the effective potential. The effective potential can be written as

$$V = \frac{\lambda}{6} \left(\Phi^\dagger \Phi - \frac{1}{2} k^2 \right)^2 \quad (4)$$

We choose the axis in such a way that $\phi_1 = \phi_2 = \phi_3 = 0$. So eq. (4) becomes

$$V = \frac{\lambda}{24} (\phi^2 - k^2)^2.$$

The mass generating term comes from the interaction term $(D^\mu \Phi)^\dagger (D_\mu \Phi)$ where the covariant derivative is

$$(D_\mu \Phi) = \left(\partial_\mu - \frac{i}{2} g \sigma \cdot W_\mu - \frac{i}{2} g' B_\mu - \frac{i}{2} g'' B''_\mu \right) \Phi, \quad (6)$$

where σ 's are the Pauli isospin matrices, $W_{\mu,i}$ are gauge bosons of $SU(2)_L$, B_μ and B''_μ are the vector bosons of the two $U(1)$'s; g, g' and g'' are the coupling constants, respectively. The ϕ -dependent mass squares of all the particles of the model can be written in a general form $m_j^2(\phi) = a_j + b_j \phi^2$ [16] as

$$m_H^2(\phi) = \frac{\partial^2 V}{\partial \phi^2} = \frac{\lambda}{6} (3\phi^2 - k^2),$$

$$m_G^2(\phi) = \frac{1}{\phi} \frac{\partial V}{\partial \phi} = \frac{\lambda}{6} (\phi^2 - k^2),$$

$$m_{W_i}^2(\phi) = \frac{m_W^2}{k^2} \phi^2$$

$$m_{B'}^2(\phi) = \frac{m_{B'}^2}{k^2} \phi^2,$$

$$m_{B''}^2(\phi) = \frac{m_{B''}^2}{k^2} \phi^2,$$

and

$$m_{Z'}^2(\phi) = \frac{m_{Z'}^2}{k^2} \phi^2,$$

where $m_G(\phi)$ is the mass function of each Goldstone boson. Quadratic divergences [18] in the Standard Model were first

studied by Veltman [19] in the context of Dimensional Regularization (DREG). He showed that in the case of regularization by Dimensional Reduction (DRED) [20] method, in the absence of quadratic divergences at one loop in the SM, the relation between the mass squares of the particles could be written as

$$12m_t^2 = 3m_H^2 + 6m_W^2 + 3m_Z^2 + 3m_{Z'}^2 \quad (13)$$

where m_H , m_W , m_Z , $m_{Z'}$ and m_t are the masses of Higgs boson, W -boson, Z -boson, Z' -boson and top quark, respectively. This equation has been derived in Ref. [21]. Eq. (13) renders the radiative corrections to the Higgs-boson mass free of quadratic divergences at one loop. From eq. (13), it is clear that the coefficients of the terms m_W^2 , m_Z^2 and $m_{Z'}^2$ are $2(d'-1)$, $(d'-1)$ and $(d'-1)$, respectively, where $d' = g_{\mu\nu} g^{\mu\nu}$ and $g_{\mu\nu}$ is the metric tensor. For DRED method, $d' = 4$; but in the conventional Dimensional Regularization (DREG) method, $d' = d$ where

$$d = 4 - \frac{\epsilon}{L}, \quad (14)$$

and L is the number of loops. Here, we consider one loop in the Standard Model, hence eq. (13) becomes

$$12m_t^2 = 3m_H^2 + 2m_W^2 + m_Z^2 + m_{Z'}^2. \quad (15)$$

This is the relation between the mass squares of the particles in DREG at one loop in the Standard Model.

3. Results and discussion

In this section, we want to calculate the mass of Z' -boson using the experimental values of the parameters in eqs. (13) and (15). The value of m_H in the analysis gives $m_H = 88_{-37}^{+60}$ GeV [22] with global fit of electroweak data [23]. However, in a supersymmetric extension of the Standard Model, there is always one light Higgs boson with mass $m_H \leq 130$ GeV in the minimal version and $m_H \leq 200$ GeV in a more general one, but the only available information on the lower limit of $m_H \geq 114.1$ GeV is established, at 95% confidence level, on the mass of the Standard Model Higgs boson at LEP [24]. However, if the SM is extended to the GUT scale $\Lambda_{GUT} \sim 10^{16}$ GeV, including top quark loops on the running coupling, the Higgs mass would roughly lie in the range of 130 GeV $\leq m_H \leq 180$ GeV [25]. In another consideration [26], it has been shown that the mass of m_H has the upper limit $m_H < 200$ GeV, whereas the cross section for $e^+e^- \rightarrow \tau\bar{\tau}HH$ [27], is consistent with the mass limit 1.5 TeV $< m_R < 1.6$ TeV. There is some evidence for the detection of Higgs boson at its mass of about 115 GeV, from electron-positron interactions at LEP [28]. Renton [29] has estimated the mass of

Higgs boson in the range of 100 - 200 GeV. Recently, we [30] have predicted the mass of Higgs boson in a flavour-independent potential model as $m_H = 120$ GeV.

We take $m_W = 80.4245 \pm 0.038$ GeV, $m_Z = 91.1876 \pm 0.0021$ GeV and $m_t = 178.1_{-8.3}^{+10.4}$ GeV [31] for our calculations. With these values, our calculations give a set of values of masses as:

$$m_{Z'} = 562.3 \pm 3.1 \text{ GeV}$$

$$\text{and } m_H = 120.0 \pm 5.0 \text{ GeV.} \quad (16)$$

Again if we take $m_H = 100$ GeV, we get $m_{Z'} = 573.9$ GeV and for $m_H = 200$ GeV, we get $m_{Z'} = 489.2$ GeV.

A broad class of supersymmetric extensions of the Standard Model predict a Z' -boson whose mass is naturally in the range of 250 GeV $< m_{Z'} < 2$ TeV [32]. There exists an experimental lower limit of about 500 GeV [33] for the Z' -gauge boson in the interaction comparable to the other couplings of the Standard Model. However, the lower mass limit can be as low as 130 GeV [34] if the coupling is weak. In the analysis of the decays of B -meson in Z' -mediated flavour-changing neutral current (FCNC) consideration [35], it has set a model-dependent lower bound of mass of Z' -boson around 500 GeV and the upper bound around 1 TeV. Carena *et al* [36] have studied the discovery potential of the Tevatron for a Z' -boson. From their study, the D-Zero detector (D0) [37] and the Collider Detector at Fermilab (CDF) [38] experiments are expected to probe Z' -boson masses in 500-800 GeV range. More interestingly, our estimation gives the mass of Z' -boson in the range of 489.2 - 573.9 GeV, which is accessible to experiments at low energies. These facts lead to enrichment in the phenomenology of Z' -bosons and physics beyond the Standard Model will show up after the discovery of the Z' -boson.

Acknowledgments

We would like to thank Prof. L Maharana, Utkal University, Bhubaneswar, for helpful discussions and suggestions. We have profited from a discussion with K Senapati, Indian Institute of Technology, Kanpur which is thankfully acknowledged. We thank the referee for suggesting valuable improvements of the manuscript.

References

- [1] M Masip and A Pamarol *Phys. Rev. D* **60** 096005 (1999); G-C Cho *Mod. Phys. Lett. A* **15** 311 (2000); K R Lynch *et al*, *Phys. Rev. D* **63** 035006 (2001); H Georgi and S L Glashow *Phys. Rev. Lett.* **32** 438 (1974); H Fritzsch and P Minkowski *Ann. Phys.* **93** 193 (1975)
- [2] M Cvetič and P Langacker *Mod. Phys. Lett. A* **11** 1247 (1996); M Cvetič, D A Demir, J R Espinosa, L Everett and P Langacker *Phys. Rev. D* **58** 119905 (1998)
- [3] J Hewett and T Rizzo *Phys. Rep.* **183** 193 (1989); A Leike *Phys. Rep.* **317** 143 (1999); M Cvetič and S Godfrey hep-ph/9504216

- [4] F Abe *et al* (CDF Collaboration) *Phys. Rev. Lett.* **67** 2418 (1991)
- [5] B B Deo and L Maharana *J. Phys. G.* **27** 815 (2001)
- [6] F Abe *et al* (CDF Collaboration) *Phys. Rev. Lett.* **79** 2192 (1997)
- [7] J Erler and P Langacker *Phys. Lett.* **B456** 68 (1999); J Erler and P Langacker *Phys. Rev. Lett.* **84** 212 (2000)
- [8] *The LEP Electroweak Working Group and SLD Heavy Flavour Group*, hep-ex/0212036
- [9] G P Zeller *et al* (NuTeV Collaboration) *Phys. Rev. Lett.* **88** 091802 (2002)
- [10] P Langacker and M Plumacher *Phys. Rev.* **D62** 013006 (2000), hep-ph/0001204
- [11] J S Ginges and V V Flambaum, arXiv:physics/0309054
- [12] L Everett, P Langacker, M Plumacher and J Wang, hep-ph/0001073
- [13] *The LEP Electroweak Working Group and the SLD Heavy Flavour Group*, Note LEPEWWG/2003-01; <http://lepewwg.web.cern.ch/LEPEWWG>
- [14] R W Robinett and J L Rosner *Phys. Rev.* **D25** 3036 (1982); M H Austern, *Ph. D. Thesis* (University of California at Berkeley, USA) (1994)
- [15] J C Pati and A Salam *Phys. Rev.* **D10** 275 (1974); R N Mohapatra and J C Pati *Phys. Rev.* **D11** 566 (1975); G Senjanovic and R N Mohapatra *Phys. Rev.* **D12** 1502 (1975); G Senjanovic *Nucl. Phys.* **B153** 334 (1979); R N Mohapatra *Unification and Supersymmetry* (New York : Springer) (1986)
- [16] B B Deo and L Maharana *Phys. Lett.* **B323** 417 (1994); B B Deo and L Maharana *Phys. Lett.* **B461** 105 (1999), B B Deo and L Maharana *Phys. Lett.* **B597** 192 (2004)
- [17] G A Kozlov hep-ph/0501154; S Coleman and E Weinberg *Phys. Rev.* **D7** 1888 (1973); D Bailin and A Love *Introduction to Gauge Field Theory* (First Indian Edition) (NewDelhi : Overseas Press India) (2004)
- [18] I Jack and D R T Jones *Nucl. Phys.* **B342** 127 (1990); M S Al-sarhi, I Jack and D R T Jones *Z Phys.* **C55** 283 (1992) and *Nucl. Phys.* **B345** 431 (1990)
- [19] M Veltman *Acta Phys. Pol.* **B12** 437 (1981)
- [20] W Siegel *Phys. Lett.* **B84** 193 (1979); D M Capper, D R T Jones and P Van Nieuwenhuizen *Nucl. Phys.* **B167** 479 (1980)
- [21] I Jack and D R T Jones *Phys. Lett.* **B234** 321 (1990), M B Einhorn and D R T Jones *Phys. Rev.* **D46** 5206 (1992)
- [22] *The LEP and SLC Electroweak Working Group*, hep-ex/0112021
- [23] A Djouadi *Pramana – J. Phys.* **60** 215 (2003)
- [24] *The LEP Higgs Working Group, Search for the Standard Model Higgs Boson at LEP*, hep-ex/0107029; hep-ex/0107030; p Kemenes *Pramana – J. Phys.* **62** 555 (2004)
- [25] T Hambye and K Riesselman *Phys. Rev.* **D55** 7255 (1997)
- [26] K Tobe and James D Wells *Phys. Rev.* **D66** 013010 (2002); J Bauer, R Clare, J Erler, S Heinemeyer, D Wackerroth, G Weiglein and D Wood, hep-ph/0111314
- [27] A Gutierrez – Rodriguez, M A Hernandez – Ruiz and O A Sanpauze *Phys. Rev.* **D67** 074018 (2003)
- [28] ALEPH, DELPHI, L3, OPAL, the LEP Higgs Working Group Search the Standard Model Higgs Boson at LEP *Phys. Lett.* **B565** 61 (2003) [hep-ex/0306033]
- [29] P Renton *Nature* **428** 141 (2004)
- [30] S Sahoo, A K Meikap and L Maharana *Mod. Phys. Lett.* **A21** 275 (2006)
- [31] S Eidelman *et al* (Particle Data Group) *Phys. Lett.* **B592** 1 (2004)
- [32] J D Lykken, hep-ph/9610218
- [33] F Abe *et al* (CDF Collaboration) in Ref [6], S Abachi *et al* (CDF Collaboration) *Phys. Lett.* **B385** 471 (1996), M Cvetic and P Langacker, hep-ph/9707451
- [34] M Cvetic and P Langacker *Mod. Phys. Lett.* **A11** 1247 (1996)
- [35] V Barger, C – W Chiang, P Langacker and H – S Lee *Phys. Lett.* **B580** 186 (2004)
- [36] M Carena, A Daleo, B A Dobrescu and T M P Tait, hep-ph/0408098.
- [37] DO Collaboration, note 4375-Conf. Search for heavy Z' Boson in the Dielectron Channel with 200 pb⁻¹ of Data (March 17 2004)
- [38] Tracey Pratt (for the CDF Collaboration), talk at the SUSY 2004 Conference (Epochal Tsukula, Tsukula, Japan) (June 2004)