



Electronic spectral studies of 2-chloro-6-methylpyridine in various solvents

B S Yadav*, Amit Kumar Yadav, Seema¹ and Subhash Chand²

Molecular Spectroscopy and Biophysics Laboratory, D.N (PG) College, Meerut-250 002,
Uttar Pradesh, India

¹Department of Chemistry, J V (PG) College Baraut, Bagpat-250 602, Uttar Pradesh, India

²Department of Chemistry, Meerut College, Meerut-250 001, Uttar Pradesh, India

Received 8 June 2006, accepted 8 February 2007

Abstract : The electronic transition bands of 2-chloro-6-methyl pyridine molecule in liquid state and the effect of α -halogen atom on the absorption intensity of the *B*-band (characteristic band of aromatic ring) have been studied. The π - π^* transition (E_2 -band) has shown the usual behavior towards the substitution and the polarity of the solvents. But the n - π^* transition (*R*-band) has evidenced a usual behavior towards the solvent polarity (i.e. the n - π^* transition has shown blue shift with respect to the n - π^* system of pyridine). The halogen atom and/or ortho para directing group (having high potential for a large inductive effect) have shown a marked influence on the sp^2 electrons of the N-atom of the ring. As a result, the *B*-band has evidenced an unusual behavior i.e. it has not shown a hyperchromic effect with increase in the polarity of the solvent.

Keywords : Ultra violet absorption spectra, solvents effect on electronic transitions

PACS Nos : 78.40. Dw, 31. 70 Dk

1. Introduction

The N-Heterocyclic compound such as Pyridine and their derivatives are of considerable biological importance. They have been known to act as herbicides, insecticides, rodenticides and plant growth regulators [1-4] etc. However very little spectroscopic studies are reported for them in literature. Pyridine shows in addition to the usual π - π^* system, a shoulder band towards longer wave lengths. This system which is symmetry allowed and polarized perpendicular to the molecular plane, has been assigned as due to n - π^* transition by Kasha [5]. Further, a systematic and complete Vibrational analysis of electronic spectra of Pyridine and some of its deuterated analogues has been made by some previous workers [6-8]. On substitution, the π - π^* and n - π^* system of Pyridine normally produce

* Corresponding Author

bathochromic and hypsochromic shifts respectively, the vibrational frequencies also get modified. As regards the earlier studies, the spectral data of some of the mono and disubstituted pyridine are provided among other [5, 9-15]. In the present work, we have measured the $\pi\text{-}\pi^*$ and $n\text{-}\pi^*$ transitions of 2-chloro-6-methyl pyridine (here after referred as 2,6-CMP) in the region of 400-200 nm in liquid state using polar solvents with a view to investigate the effect of solvent and effect of halogen and/or ortho para directing groups on the position and intensity of the electronic absorption bands.

2. Experimental procedure

The spec-pure grade sample of 2, 6-CMP was obtained from M/s Sigma Aldrich Chemicals Co. West Germany and used as such. The Purity of the compound was further confirmed by boiling point (64°) determination and elemental analysis. We have measured the $n\text{-}\pi^*I$, $n\text{-}\pi^*II$ and $\pi\text{-}\pi^*$ transitions of 2-chloro-6-methyl pyridine in a uv/vis double beam spectrophotometer Model Backman M-35 in the region of 400-200 nm liquid state, using various polar solvents (*viz*-ethanol→methanol→water). The solvent used for preparing the solutions of the said compound were of spectroscopic grade.

3. Results and discussion

The molecular structure of 2, 6-CMP is shown in Figure 1. The near ultra violet absorption spectra (region 400-200nm) of 2, 6-CMP in various solvents (a) Ethanol (b) Methanol and (c) Water are shown in Figure 2. The electronic transitions of the compound observed in various solvents are given in Table 2.




Figure 1. The molecular structure of 2-chloro-6-methyl pyridine.

3.1 Band due to $\pi\text{-}\pi^*$ transition :

The spectra of heteroaromatic compounds are on the whole similar to their corresponding hydrocarbons[16,17]. The spectrum of pyridine is comparable with that of benzene. Benzene itself displays three electronic absorption bands[16,17] at 184nm (ϵ , the molar absorptivity or the extinction = 60,000), at 203.5nm ($\epsilon=7900$) and at 256nm ($\epsilon=200$) due to $\pi\text{-}\pi^*$ transitions. The intense band at 180nm results from an allowed transition[17], whereas the weaker bands at 200 and 260nm result from forbidden transitions in the highly symmetrical benzene molecule. Different notations have been used to designate the absorption bands of benzene, these are summarized in Table 1. We shall discuss these




Figure 2. Near ultra violet absorption spectra (region 400-200 nm) of 2-chloro-6-methyl pyridine in (a) ethanol (b) methanol and (c) water.

using braude's *E* and *B* notations[17]. The *B*-band has evidenced an unusual behavior *i.e.* it has not shown a hyperchromic effect with increase in the polarity of the solvent. Such type of bands *i.e.* unusual behaviors of *B*-band arises due to π - π^* transition in aromatic or hetro-aromatic molecules (the π - π^* transition has shown red shift with respect to the π - π^* system of pyridine). Benzenes shows absorption peaks between 230-270 nm. When a chromophoric group is attached to the benzene ring, the *B*-bands are observed at longer

wave-lengths than the more intense *K*-bands. The position as well as the intensity of the *B*-band is not shifted by increasing the polarity of the solvent. But in heterocyclic aromatic compounds a marked hyperchromic shift (increases in ϵ_{max}) is observed by increasing the polarity of the solvent. This type of transition occurs in the unsaturated centers of the molecule, *i.e.*, in compounds containing double or triple bonds and also in aromatics. The excitation of π electron requires smaller energy and hence, transition of this type occurs at longer wavelength. A π electron of a double bond is excited to π^* orbital. *B*-band is generally found in between 230-270 nm.

Table 1. Electronic absorption bands of benzene molecule (16).

184nm	204nm	256nm	References
E_1 -and	E_2 -and	<i>B</i> -band	(18)
-	<i>K</i> -band	<i>K</i> -band	(19)

Between benzene and pyridine, the *B*-band of pyridine is more intense with some what diminished fine structure where as in benzene, the *B*-band at 256nm displays fine structure *i.e.* the band contains multiple peaks. This transition is allowed for pyridine but forbidden for more symmetrical benzene [16,17]. The band observed at 282nm, 283nm and 277nm in the study in all the three solvents (ethanol, methanol and water) are originated from π - π^* transition and designated as *B*-band.

3.2 Bands due to n - π^* transition :

The *R*-band expected for an n - π^* transition for pyridine molecule has been observed in vapour phase spectra. The spectrum is determined in solution [17] because of intermolecular hydrogen bonding found between the N-atom of pyridine ring and the solvent molecule.

The *R*-bands found in this study, have originated from n - π^*I and n - π^*II transitions due to the attachment of groups containing lone pair electrons *i.e.* -Cl or -CH₃ groups and occurred at 298nm (Methanol), 291nm (Ethanol/Methanol) and 281nm (Water) respectively.

4. Effect of substitution on electronic transitions

An Auxochrome (*i.e.* a saturated group with non bonded electrons *eg*: -OR, -ON, -NH₂, and -Cl *etc.*) attached to a chromophore (a covalently unsaturated group responsible for electronic absorption *eg*. aromatic ring, -C=C-, -C≡C-, *etc.*) alters both the wave length and the intensity of the absorption [16]. Auxochromic substitution on the aromatic ring shifts *B*-band to longer wave length, frequently with intensification of the *B*-band and loss of its fine structure because of π - π^* conjugation [17]. Generally, the *B*-band occurred in between (230-270nm) and displays a fine structure *i.e.* the band contain multiple peaks. In the present investigation *B*-bands (π - π^*) observed at 282nm, 283nm and 277nm in ethanol→methanol→water have large intensity without a fine structure. In auxochromic substitution. The *B*-bands shifts to longer wavelength frequently increase in intensity.




Figure 3. The combined inductive and mesomeric (resonance) effects of chloro and methyl group responsible for shift in $n-\pi^*$ and $\pi-\pi^*$ transitions.

The auxochromic groups present on aromatic ring are responsible for $n-\pi^*$ transition (R -band) due to the presence of non bonding electrons (*i.e.* lone pair electrons), available on the auxochromic group. In 2,6-CMP molecule (*i.e.* disubstituted molecule), the inductive effect of auxochromic groups is responsible for $n-\pi$ conjugation. The -Cl group has three lone pair of electrons to that of $-\text{CH}_3$ group. Further more, the -Cl group is highly electronegative and have much more negative inductive effect in comparison to that of $-\text{CH}_3$ group and the size of -Cl group is much lower than that of $-\text{CH}_3$ group. As a result of which valence shell electrons of -Cl group contribute large nuclear charge in comparison to that of $-\text{CH}_3$ group. In conjugated systems, the energy separation between the groups at the excited states is reduced and the system absorbs at larger wave lengths with an increased intensity [16]. (*i.e.* B -band is intense at longer wave length). The shift in absorption results from a combination of inductive and resonance effects are shown in Figure 3. The R -band (due to the additive effect of auxochromic substitutions) shifts to longer wave lengths at 298nm (Methanol), 291nm (Ethanol/Methanol) and 281nm (Water) respectively.

5. Effect of solvents on electronic transitions

The B -band of Benzene, an increase in the solvent polarity does not effect either the position or the intensity. But Produces a hyperchromic effect (an increase in absorption intensity). For pyridine and its homologs [16,17], this effect is assigned to the hydrogen bonding through the lone pair of electrons of the nitrogen atom [18,19] In 2,6-CMP molecule, an increase in solvent polarity, hyperchromic effect is not observed because of the presence of α -halogen atom (-Cl), which has high negative inductive effect (-I) and influence that sp^2 electrons of the N -atom of the ring, As a result, it makes the lone pair electrons of N -atom of the ring less available for H -bonding. The negative inductive effect of chloro group becomes much greater, as the solvent polarity increases by the dipole-dipole interaction between the solute and solvent particles. As the solvent polarity decreases, the negative inductive effect of -Cl group also decreases. As a result, the lone pair electrons of N -atom of the ring are readily available for intermolecular H -bonding to result in the increase in absorption intensity of the B -band in less polar solvent than in more polar solvent.

The $n-\pi^*$ transition (R -band) shows a hypsochromic shift (blue shift) on increasing the polarity of the solvent [16]. The $n-\pi^*$ transition will be of higher energy *i.e.* of lower wave length (blue shift) which is shown in Figure 4. In the present work the blue-shift has been observed by $n-\pi^*$ transition (R -band) as the polarity of the solvent increased [20,21] *i.e.*

ethanol→methanol→water (Table 2). The amount of blue-shift has been used as a measure of the strength of the *H*-bonding.




Figure 4. Solvent effect on $n-\pi^*$ and $\pi-\pi^*$ transition.

The $\pi-\pi^*$ transition has shifted to the longer wave length *i.e.* red shifted (Lower energy) in a more polar solvent [16]. The polar excited state of the $\pi-\pi^*$ transition is stabilized by *H*-bonding in more polar solvent. Becker *et al* [22], has suggested that the $\pi-\pi^*$ transition under go red shifts (Lower energy) as the polarity of the solvent increases. This Lowers the distance between π and π^* levels with a subsequent lowering in the energy or raising the wave length of the transition shown in Figure 4. We have observed the red shift by the $\pi-\pi^*$ transition (E_2 -band) as the polarity of the solvent has increased from ethanol→methanol→water (Table 2) in 2, 6-CMP.

Table 2. Effect of solvents on electronic absorption bands/transition of 2-chloro-6-methyl pyridine (all values are in nanometer).

Solvent	$n-\pi^*$ I (<i>R</i> -band)/nm	$n-\pi^*$ II (<i>R</i> -band)/nm	$\pi-\pi^*$ (<i>B</i> -band)/nm
Ethanol	-	291	282
Methanol	298	291	283
Water	-	281	277

Acknowledgment

Authors are thankful to Punjab University, Chandigarh for providing the necessary facility for recording the ultraviolet absorption spectra.

References

- [1] W Hoyle and G A Howarth *Chem. Abstr.* **78** 1608h (1973)
- [2] Tokeda Chem. Industries Ltd., *Jpn. Chem. Abstr.* **80** 89650b (1974)
- [3] B S Yadav, Seema and Vipin Kumar *Indian J. Pure Appl. Phys* **35** 587(1997)
- [4] B S Yadav, M K Yadav and Vipin Kumar *Acta Botanica Indica* **26** 11(1998)
- [5] M Kasha *Faraday Disc. Soc.* **9** C14 (1954)
- [6] K C Medhi *Spectrochim. Acta* **38A** 717 (1982)
- [7] B S Yadav, Nitin Kumar and Seema *Ultra Science* **17(3)** 469 (2005)
- [8] R V Shanbhag M A Shashidhar and K Surya Narayan Rao *Indian J. Phys.* **58B** 248 (1979)
- [9] B S Yadav, Karuan Gupta and Subhas Chand *Indian J. Phys.* **77B (4)** 417 (2003)
- [10] T N Misra *Indian J. Phys.* **35** 420 (1961)
- [11] B R Pandey and R S Tripathi *Indian J. Pure Appl. Phys.* **12** 64 (1974)
- [12] B S Yadav, Nitin Kumar and Vipin Kumar *Orient J. Chem*, **21(2)** 255 (2005)
- [13] H P Stephersson *J. Chem. Phys.* **22** 1077 (1954)
- [14] H Sponer and J H Rushi *J. Chem. Phys.* **20** 1847 (1952)
- [15] S L Shrivastava and M Prasad *Spectrochim. Acta.* **40A** 1101 (1984)
- [16] P S Kalsi *Spectroscopy of Organic Compounds* (New Delhi : New Age International) Ch 2. p.7 (2001)
- [17] R M Silverstein, G C Bassler and T C Morrill *Spectrometric Identification of Organic Compounds* (New York : John Wiley) Ch 6, p.305 (1981)
- [18] E A Braude *U V and Visible Light Absorption in Determination of Organic Structure by Physical Methods* (New York : Academic) Ch 4, p94 (1955)
- [19] J Buraway *Chem. Soc.* **63** 3155 (1930)
- [20] B S Yadav and V Singh *Spectrochim. Acta.* **55A** 1267 (1999)
- [21] K Ram, B R Pandey and R S Tripathi *J. Chem. Phys.* **74** 1250 (1977)
- [22] R S Becker, A B F Duncan, F A Matsen, DR Scoot and W West *Chemical Application of Spectroscopy* (New York : John Wiley) (1968)